

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

HIGH FIDELITY DNA POLYMERASE COMPOSITIONS AND USES THEREFOR

This application is a continuation in part of U.S. Patent application with serial No.
10 10/035,091, filed December 21, 2001.

FIELD OF THE INVENTION

The present invention is related to the field of high fidelity polynucleotide synthesis.

BACKGROUND OF THE INVENTION

DNA polymerases catalyze the synthesis of DNA and can be found in all cells as well as
being encoded in numerous viruses. Although all DNA polymerases possess 5'-3' DNA
polymerization activity, DNA polymerases differ from one another in numerous other properties.
For example, some enzymatic activities that are possessed by some DNA polymerases, but
absent in other DNA polymerases include: double stranded DNA 5'-3' exonuclease activity,
single-stranded DNA 3'-5' exonuclease activity, double-stranded 3'-5' DNA exonuclease activity,
RNase H activity, reverse transcriptase activity, and the like. Additionally, different DNA
polymerases may have different optimal pH and temperature ranges for activity. Furthermore,
DNA polymerases may differ in the rate in which they catalyze DNA synthesis.

Purified DNA polymerases have numerous uses in vitro. A detailed description of DNA
polymerases, including methods for their isolation, can be found among other places, in DNA
25 Replication 2nd edition, by Kornberg and Baker, W. H. Freeman & Company, New York, N.Y.
1991. In vitro uses of DNA polymerases include, for example, the labeling and synthesis of
hybridization probes, DNA sequencing, and DNA amplification. A DNA amplification method
employing DNA polymerases that has been particularly useful is the polymerase chain reaction
(PCR) technique which employs the use of a thermostable DNA polymerase.

30 The first thermostable DNA polymerase that is widely used for DNA amplification is Taq
DNA polymerase isolated from the thermostable, aerobic bacterium *Thermus aquaticus*. Taq
DNA polymerase's enzymatic activity at high temperatures allows for primer extension and

5 sequencing of polynucleotide templates with complex secondary structures (i.e., by PCR amplification). However, Taq DNA polymerase has significant error rate when incorporating nucleotides due to the lack of 3'-5' exonuclease activity (i.e., proofreading activity), and therefore may not be suitable if the amplified sequence is to be used in further gene structural/functional studies or cloning.

10 In the last 10 years, numerous studies have quantified the error rate of thermostable DNA polymerases, and several enzymes have been found to copy DNA more accurately than Taq DNA polymerase (referred to as high fidelity DNA polymerases). U.S. Patent describing DNA polymerases include Nos. 4,492,130; 4,946,786; 5,210,036; 5,420,029; 5,489,523; 5,506,137; 5,545,552; 5,618,711; 5,624,833; 6,238,905; 6,100,078; 6,077,664; 5,968,799; 5,948,663; 5,885,713; 5,834,285; 5,756,334; 5,747,298; 5,744,312; 5,624,833; 5,602,011; 5,556,772.

15 High fidelity polymerases alone should definitely increase fidelity rates but usually do not amplify long fragments as efficient as a DNA polymerase lacking a 3'-5' exonuclease activity (e.g., Taq DNA polymerase). Enzyme mixtures that combine a standard polymerase with a small amount of proofreading polymerase may provide a balance between fidelity and yield. A study published in 1994 illustrated that the use of a high level of a DNA polymerase lacking 3'-5' exonuclease activity (an exo^- DNA polymerase, KlenTaq-1) with a very low level of a thermostable DNA polymerase exhibiting 3'-5' exonuclease activity (an exo^+ DNA polymerase such as Pfu, Vent, or Deep Vent) generated products with increased base-pair fidelity with a maximum yield of 35 kb DNA from 1 ng of lambda DNA template (Barnes, Proceedings of the
20 National Academy of Sciences, 91:2216-20, 1994). Similarly, U.S. Patent Nos. 5,436,149 and 6,008,205 disclosed methods for improving DNA amplification fidelity using a DNA polymerase composition comprising a first enzyme substantially lacking 3'-5' exonuclease activity and a second enzyme comprising 3'-5' exonuclease activity. In mixtures such as these, the exo^+ enzyme acts to correct polymerization errors produced by the exo^- DNA polymerase.

25 The problem inherited in the above composition comprising the mix of two DNA polymerases is that the high polymerization activity resulted from combining the two DNA polymerases may inhibit the efficiency and therefore the yield of the amplification reaction. Therefore, one can not increase fidelity by increasing the proportion of the proofreading DNA

5 polymerase without compromising PCR product yield. It is also known that the amplification fidelity may also be affected by high DNA polymerase concentration (see for example, Mattila et al., 1991, Polynucleotides Research, 19:4967-73).

There is therefore a need in the art for new methods and compositions which improve polymerization fidelity and reduce the side effects resulted from having high polymerization
10 activity in the reaction.

SUMMARY OF THE INVENTION

The invention provides an enzyme mixture for DNA synthesis comprising a first enzyme and a second enzyme, where the first enzyme comprises a DNA polymerization activity, and the second enzyme comprises a 3'-5' exonuclease activity and a reduced DNA polymerization activity.
15

In one embodiment, the first enzyme is a DNA polymerase or a reverse transcriptase.

Preferably, the DNA polymerase is selected from the group consisting of: Taq DNA polymerase, Tth DNA polymerase, UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D
20 DNA polymerase and DP1/DP2 DNA polymerase.

Also preferably, the second enzyme is a mutant DNA polymerase.

In one embodiment, the mutant DNA polymerase is derived from a DNA polymerase different from the first enzyme.

The invention also provides an enzyme mixture for DNA synthesis comprising a first
25 enzyme and a second enzyme, where the first enzyme is a wild type Pfu DNA polymerase, the second enzyme is a mutant Pfu DNA polymerase comprising a 3'-5' exonuclease activity and a reduced DNA polymerization activity.

The invention further provides an enzyme mixture for DNA synthesis comprising a first enzyme and a second enzyme, where the first enzyme is a Taq DNA polymerase, the second

5 enzyme is a mutant Pfu DNA polymerase comprising a 3'-5' exonuclease activity and a reduced DNA polymerization activity.

Preferably, the mutant DNA polymerase is derived from a DNA polymerase selected from the group consisting of: UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D
10 DNA polymerase and DP1/DP2 DNA polymerase.

Also preferably, the mutant Pfu DNA polymerase comprises one or more mutations at amino acid positions selected from the group consisting of: D405, Y410, T542, D543, K593, Y595, Y385, G387, and G388.

In some embodiments, the mutant DNA polymerase is one selected from the group of
15 mutations in Table 2A and 2B.

Still more preferably, the mutant Pfu DNA polymerase comprises one or more mutations selected from the group consisting of: D405E, Y410F, T542P, D543G, K593T, Y595S, Y385Q, Y385S, Y385N, Y385L, Y385H, G387S, G387P, and G388P.

The enzyme mixture of the present invention may further comprise a PCR enhancing
20 factor and/or an additive.

The invention provides a kit for DNA synthesis comprising a first enzyme and a second enzyme, where the first enzyme comprises a DNA polymerization activity, the second enzyme comprises a 3'-5' exonuclease activity and a reduced DNA polymerization activity, and packaging material therefore.

25 Preferably, the first enzyme in the kit is a DNA polymerase or a reverse transcriptase.

More preferably, the DNA polymerase in the kit is selected from the group consisting of: Taq DNA polymerase, Tth DNA polymerase, UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D DNA polymerase and DP1/DP2 DNA polymerase.

30 Preferably, the second enzyme in the kit is a mutant DNA polymerase.

5 More preferably, the mutant DNA polymerase is derived from a DNA polymerase selected from the group consisting of: UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D DNA polymerase and DP1/DP2 DNA polymerase.

10 Still more preferably, the mutant DNA polymerase is derived from a DNA polymerase different from the first enzyme.

The invention provides a kit comprising an enzyme mixture for DNA synthesis, the kit comprises a first enzyme and a second enzyme, and packaging material therefore, where the first enzyme is a wild type Pfu DNA polymerase, the second enzyme is a mutant Pfu DNA polymerase comprising a 3'-5' exonuclease activity and a reduced DNA polymerization activity.

15 The invention also provides a kit comprising an enzyme mixture for DNA synthesis, the kit comprises a first enzyme and a second enzyme, and packaging material therefore, where the first enzyme is a Taq DNA polymerase, and packaging material therefore, the second enzyme is a mutant Pfu DNA polymerase comprising a 3'-5' exonuclease activity and a reduced DNA polymerization activity.

20 The kit of the invention may further comprises one or more components selected from the group consisting of: a deoxynucleotide, a reaction buffer, a PCR enhancing factor and/or additive, a control DNA template and a control primer.

25 The mutant Pfu DNA polymerase of the kit may comprise one or more mutations at amino acid positions selected from the group consisting of: D405, Y410, T542, D543, K593, Y595, Y385, G387, and G388.

Preferably, the the mutant Pfu DNA polymerase comprises one or more mutations selected from the group consisting of: D405E, Y410F, T542P, D543G, K593T, Y595S, Y385Q, Y385S, Y385N, Y385L, Y385H, G387S, G387P, and G388P.

The invention provides a method for DNA synthesis comprising:

5 (a) providing an enzyme mixture, the enzyme mixture comprising a first enzyme comprising a DNA polymerization activity, and a second enzyme comprising a 3'-5' exonuclease activity and a reduced DNA polymerization activity; and

(b) contacting the enzyme mixture with a nucleic acid template, where the enzyme mixture permits DNA synthesis.

10 Preferably, in the method of the invention, the nucleic acid template is a DNA or an RNA molecule.

Also preferably, the first enzyme used in the method is a DNA polymerase or a reverse transcriptase.

15 More preferably, the DNA polymerase is selected from the group consisting of: Taq DNA polymerase, Tth DNA polymerase, UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D DNA polymerase and DP1/DP2 DNA polymerase.

Preferably, the mutant DNA polymerase comprises a mutation in its partitioning domain or the polymerase domain.

20 Also preferably, the second enzyme is a mutant DNA polymerase.

More preferably, the mutant DNA polymerase is derived from a DNA polymerase selected from the group consisting of: UITma DNA polymerase, Tli DNA polymerase, Pfu DNA polymerase, KOD DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase, PGB-D DNA polymerase and DP1/DP2 DNA polymerase.

25 Still more preferably, the mutant DNA polymerase is derived from a DNA polymerase different from the first enzyme.

The invention also provides a method for DNA synthesis comprising:

5 (a) providing an enzyme mixture, the enzyme mixture comprising a wild type Pfu DNA polymerase as a first enzyme, and a mutant Pfu DNA polymerase as a second enzyme which comprises a 3'-5' exonuclease activity and a reduced DNA polymerization activity; and

(b) contacting the enzyme mixture with a nucleic acid template, where the enzyme mixture permits DNA synthesis.

10 The invention further provides a method for TA cloning of DNA synthesis product comprising:

(a) providing an enzyme mixture, the enzyme mixture comprising a Taq DNA polymerase as a first enzyme, and a mutant Pfu DNA polymerase as a second enzyme which comprises a 3'-5' exonuclease activity and a reduced DNA polymerization activity;

15 (b) contacting the enzyme mixture with a nucleic acid template, where the enzyme mixture permits DNA synthesis to generate a synthesized DNA product; and

(c) inserting the synthesized DNA product into a TA cloning vector.

20 Preferably, the mutant Pfu DNA polymerase used in the method of the invention comprises one or more mutations at amino acid positions selected from the group consisting of: D405, Y410, T542, D543, K593, Y595, Y385, G387, and G388.

More preferably, the mutant Pfu DNA polymerase comprises one or more mutations selected from the group consisting of: D405E, Y410F, T542P, D543G, K593T, Y595S, Y385Q, Y385S, Y385N, Y385L, Y385H, G387S, G387P, and G388P.

25 The reaction mixture of the subject method may further comprise a PCR enhancing factor and/or an additive.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1. PCR proofreading activity assay using Pfu DNA polymerase mutants according to some embodiments of the invention.

5 Figure 2. PCR performance of Pfu plus Pfu G387P mutant blends according to some
embodiments of the invention.

 Figure 3. PCR performance of Taq plus Pfu G387P mutant blends according to
some embodiments of the invention.

DETAILED DESCRIPTION OF THE INVENTION

10 The subject invention provides novel composition for high fidelity polynucleotide
synthesis, particularly DNA synthesis. The subject compositions comprise an enzyme mixture
for DNA synthesis comprising a first enzyme and a second enzyme, where the first enzyme
comprises a DNA polymerization activity, and the second enzyme comprises a 3'-5' exonuclease
activity and a reduced DNA polymerization activity. In addition to providing high fidelity for
15 DNA synthesis, the compositions of the subject invention prevent side effects of a high
polymerization activity, therefore, increase the efficiency of the amplification compared to a
mixture in which both DNA polymerases possess wild-type polymerization activities.

Definitions

20 As used herein, "synthesis" refers to any in vitro method for making new strand of
polynucleotide or elongating existing polynucleotide (i.e., DNA or RNA). Synthesis, according
to the invention, include amplification, which increases the number of copies of a polynucleotide
template sequence with the use of a polymerase. Polynucleotide synthesis (e.g., amplification)
results in the incorporation of nucleotides into a polynucleotide (i.e., a primer), thereby forming a
new polynucleotide molecule complementary to the polynucleotide template. The formed
25 polynucleotide molecule and its template can be used as templates to synthesize additional
polynucleotide molecules.

 "DNA synthesis", according to the invention, includes, but are not limited to PCR,
reverse transcription, the labelling of polynucleotide (i.e., for probes and oligonucleotide
primers), polynucleotide sequencing.

30 As used herein, the term "template dependent manner" is intended to refer to a process
that involves the template dependent extension of a primer molecule (e.g., DNA synthesis by

5 DNA polymerase). The term “template dependent manner” refers to polynucleotide synthesis of RNA or DNA wherein the sequence of the newly synthesized strand of polynucleotide is dictated by the well-known rules of complementary base pairing (see, for example, Watson, J. D. et al., In: Molecular Biology of the Gene, 4th Ed., W. A. Benjamin, Inc., Menlo Park, Calif. (1987)).

10 As used herein, “polynucleotide polymerase” refers to an enzyme that catalyzes the polymerization of nucleotide (i.e., the polymerase activity). Generally, the enzyme will initiate synthesis at the 3'-end of the primer annealed to a polynucleotide template sequence, and will proceed toward the 5' end of the template strand. “DNA polymerase” catalyzes the polymerization of deoxynucleotides.

15 As used herein, the “polymerase domain” refers to the one or more domains of a DNA polymerase which is critical for its polymerization activity. The position of the polymerase domain varies, for example, the polymerase domain for Pfu, Tgo, KDO, Tli (Vent) and PGB-D (dee Vent) are located at amino acid positions as described in Table 2B.

20 As used herein, the “partitioning domain” refers to a domain of a DNA polymerase which plays a critical role in coordinating the balance between synthesis and degradation of the DNA chain. Generally the partitioning domain is characterized by the YXGG motif (Truniger et al., 1996, EMBO J. 15:3430-3441). This region is located within an accessible loop connecting the 3'-5' exonuclease and polymerase domains. The position of the partitioning domain varies. For example, the partitioning domain for Pfu, Tgo, KDO, Tli (Vent) and PGB-D (dee Vent) are located at amino acid positions 384-389, 383-388, 383-388, 386-391, and 384-389 respectively.

25 According to the invention, another class of DNA polymerase is “reverse transcriptases”, also referred to as “RT”, is a critical enzyme responsible for the synthesis of cDNA from viral RNA for all retroviruses, including HIV, HTLV-I, HTLV-II, FeLV, FIV, SIV, AMV, MMTV, and MoMuLV. For review, see e.g. Levin, 1997, Cell, 88:5-8; Brosius et al., 1995, Virus Genes 11:163-79. The term “reverse transcriptase (RT) activity” means the ability to synthesize cDNA
30 from RNA template. Methods for measuring RT activity are well known in the art, for example, the Quan-T-RT assay system is commercially available from Amersham (Arlington Heights, Ill.) and is described in Bosworth, et al., Nature 1989, 341:167-168.

5 As used herein, a mutant DNA polymerase with “reduced polymerization activity” is a DNA polymerase mutant comprising a DNA polymerization activity which is lower than that of the wild-type enzyme, e.g., comprising less than 10% DNA (e.g., less than 8%, 6%, 4%, 2% or less than 1%) polymerization activity of that of the wild-type enzyme.

10 As used herein, “exonuclease” refers to an enzyme that cleaves bonds, preferably phosphodiester bonds, between nucleotides one at a time from the end of a DNA molecule. An exonuclease can be specific for the 5' or 3' end of a DNA molecule, and is referred to herein as a 5' to 3' exonuclease or a 3' to 5' exonuclease. A useful exonuclease according to the invention is a 3' to 5' exonuclease which degrades DNA by cleaving successive nucleotides from the 3' end of the polynucleotide. During the synthesis or amplification of a polynucleotide template, a DNA polymerase with 3' to 5' exonuclease activity (exo^+) has the capacity of removing mispaired base (proofreading activity), therefore is less error-prone than a DNA polymerase without 3' to 5' exonuclease activity (exo^-). The exonuclease activity can be defined by methods well known in the art. For example, one unit of exonuclease activity may refer to the amount of enzyme required to cleave 1 μg DNA target in an hour at 37°C. Wild type Tth DNA polymerase and Taq DNA polymerase are “ exo^- ” because they do not have 3' to 5' exonuclease activities, however, wild type Pfu DNA polymerase, *E. coli* DNA polymerase I, T7 DNA polymerase, Tma DNA polymerase, Tli DNA polymerase, KOD DNA polymerase, JDF DNA polymerase, and PGB-D DNA polymerase are “ exo^+ ” because they all exhibit 3' to 5' exonuclease activity.

25 The term “fidelity” as used herein refers to the accuracy of DNA polymerization by template-dependent DNA polymerase. The fidelity of a DNA polymerase is measured by the error rate (the frequency of incorporating an inaccurate nucleotide, i.e., a nucleotide that is not incorporated at a template-dependent manner). The accuracy or fidelity of DNA polymerization is maintained by both the polymerase activity and the 3'-5' exonuclease activity of a DNA polymerase. The term “high fidelity” refers to an error rate of 5×10^{-6} per base pair or lower. 30 The fidelity or error rate of a DNA polymerase may be measured using assays known to the art (see for example, Lundburg et al., 1991 Gene, 108:1-6).

As used herein, an “amplified product” refers to the double strand polynucleotide population at the end of a PCR amplification reaction. The amplified product contains the

5 original polynucleotide template and polynucleotide synthesized by DNA polymerase using the polynucleotide template during the PCR reaction.

As used herein, "polynucleotide template" or "target polynucleotide template" refers to a polynucleotide containing an amplified region. The "amplified region," as used herein, is a region of a polynucleotide that is to be either synthesized by reverse transcription or amplified by
10 polymerase chain reaction (PCR). For example, an amplified region of a polynucleotide template resides between two sequences to which two PCR primers are complementary to.

As used herein, the term "primer" refers to a single stranded DNA or RNA molecule that can hybridize to a polynucleotide template and prime enzymatic synthesis of a second polynucleotide strand. A primer useful according to the invention is between 10 to 100
15 nucleotides in length, preferably 17-50 nucleotides in length and more preferably 17-45 nucleotides in length.

"Complementary" refers to the broad concept of sequence complementarity between regions of two polynucleotide strands or between two nucleotides through base-pairing. It is known that an adenine nucleotide is capable of forming specific hydrogen bonds ("base pairing")
20 with a nucleotide which is thymine or uracil. Similarly, it is known that a cytosine nucleotide is capable of base pairing with a guanine nucleotide.

The term "wild-type" refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays altered characteristics
25 when compared to the wild-type gene or gene product. For example, a mutant DNA polymerase in the present invention is a DNA polymerase which exhibit a reduced DNA polymerization activity.

As used herein, an "enzyme mixture" according to the invention, comprises a first enzyme comprising DNA polymerization activity and a second enzyme comprising a 3'-5'
30 exonuclease activity and a reduced DNA polymerization activity. The ratio of the DNA polymerase activity and the exonuclease activity in the enzyme mixture is about (2.5-5U of DNA polymerization activity)/(0.05-10U of 3'-5' exonulcease activity).

5 Useful DNA Polymerases And Reverse Transcriptases

DNA polymerases and their properties are described in detail in, among other places, DNA Replication 2nd edition, Kornberg and Baker, W. H. Freeman, New York, N.Y. (1991).

Known conventional DNA polymerases include, for example, *Pyrococcus furiosus* (Pfu) DNA polymerase (Lundberg et al., 1991, *Gene*, 108:1, provided by Stratagene), *Pyrococcus*
10 *woesei* (Pwo) DNA polymerase (Hinnisdaels et al., 1996, *Biotechniques*, 20:186-8, provided by Boehringer Mannheim), *Thermus thermophilus* (Tth) DNA polymerase (Myers and Gelfand 1991, *Biochemistry* 30:7661), *Bacillus stearothermophilus* DNA polymerase (Stenesh and McGowan, 1977, *Biochim Biophys Acta* 475:32), *Thermococcus litoralis* (Tli) DNA polymerase (also referred to as Vent DNA polymerase, Cariello et al., 1991, *Polynucleotides Res*, 19: 4193, provided by New England Biolabs), 9°Nm DNA polymerase (discontinued product from New England Biolabs), *Thermotoga maritima* (Tma) DNA polymerase (Diaz and Sabino, 1998 *Braz*
15 *J. Med. Res*, 31:1239), *Thermus aquaticus* (Taq) DNA polymerase (Chien et al., 1976, *J. Bacteriol*, 127: 1550), *Pyrococcus kodakaraensis* KOD DNA polymerase (Takagi et al., 1997, *Appl. Environ. Microbiol.* 63:4504), JDF-3 DNA polymerase (from *thermococcus sp.* JDF-3, Patent application WO 0132887), *Pyrococcus GB-D* (PGB-D) DNA polymerase (also referred as Deep-Vent DNA polymerase, Juncosa-Ginesta et al., 1994, *Biotechniques*, 16:820, provided by New England Biolabs), UITma DNA polymerase (from *thermophile Thermotoga maritima*; Diaz and Sabino, 1998 *Braz J. Med. Res*, 31:1239; provided by PE Applied Biosystems), Tgo DNA polymerase (from *thermococcus gorgonarius*, provided by Roche Molecular Biochemicals), *E.*
20 *coli* DNA polymerase I (Lecomte and Doubleday, 1983, *Polynucleotides Res.* 11:7505), T7 DNA polymerase (Nordstrom et al., 1981, *J. Biol. Chem.* 256:3112), and archaeal DP1/DP2 DNA polymerase II (Cann et al., 1998, *Proc Natl Acad Sci U S A* 95:14250-5). The polymerization activity of any of the above enzymes can be defined by means well known in the art. One unit of DNA polymerization activity of conventional DNA polymerase, according to the subject
25 invention, is defined as the amount of enzyme which catalyzes the incorporation of 10 nmoles of total deoxynucleotides (dNTPs) into polymeric form in 30 minutes at optimal temperature (e.g., 72°C for Pfu DNA polymerase). Assays for DNA polymerase activity and 3'-5' exonuclease activity can be found in DNA Replication 2nd Ed., Kornberg and Baker, supra; Enzymes, Dixon

5 and Webb, Academic Press, San Diego, Calif. (1979), as well as other publications available to the person of ordinary skill in the art.

When using the subject compositions in reaction mixtures that are exposed to elevated temperatures, e.g., during the PCR technique, use of thermostable DNA polymerases is preferred.

10 Reverse transcriptases useful according to the invention include, but are not limited to, reverse transcriptases from HIV, HTLV-1, HTLV-II, FeLV, FIV, SIV, AMV, MMTV, MoMuLV and other retroviruses (for reviews, see for example, Levin, 1997, Cell, 88:5-8; Verma, 1977, Biochim Biophys Acta. 473:1-38; Wu et al., 1975, CRC Crit Rev Biochem. 3:289-347).

15 Useful First Enzyme Comprising DNA Polymerization Activity

Enzymes comprising DNA polymerization activity according to the present invention include enzymes such as DNA polymerases and reverse transcriptases.

20 The first enzyme used in the subject composition can be any DNA polymerase, with or without a proof reading activity. Preferably, a wild type DNA polymerase is used. However, a mutant DNA polymerase can also be used so long as it provides sufficient DNA polymerization activity required for an amplification reaction.

In a preferred embodiment, the first enzyme comprising DNA polymerization activity is a wild type Pfu DNA polymerase. The enzyme mixture comprising a Pfu DNA polymerase as the first enzyme is also referred to as a Pfu blend herein after.

25 In preferred embodiments of the invention, a Pfu blend enzyme mixture is used for DNA synthesis reaction, e.g., PCR reaction.

In another preferred embodiment, the first enzyme comprising DNA polymerization activity is a wild type Taq DNA polymerase. The enzyme mixture comprising a Taq DNA polymerase as the first enzyme is also referred to as a Taq blend herein after.

5 In preferred embodiments of the invention, a Taq blend enzyme mixture is used for DNA synthesis reaction and for subsequent direct cloning, e.g., PCR reaction followed by TA cloning.

Useful Second Enzyme Comprising 3'-5' Exonuclease Activity

Enzyme comprising 3'-5' exonuclease activity (i.e., proofreading DNA polymerase) according to the invention include, but are not limited to, DNA polymerases, *E. coli* exonuclease I, *E. coli* exonuclease III, *E. coli* recBCD nuclease, mung bean nuclease, and the like (see for example, Kuo, 1994, Ann N Y Acad Sci., 726:223-34).

Any proofreading DNA polymerase could be mutagenized to reduce/eliminate DNA polymerase activity and used in the enzyme reaction of the present invention. Examples can be found in many DNA polymerase families including, but are not limited to such as follows:

Family B DNA polymerases

Bacteriophage T4 DNA polymerase, ϕ 29 DNA polymerase, T7 DNA polymerase; *E. coli* pol II DNA polymerase; human DNA polymerase δ , human DNA polymerase γ , archaeal DNA polymerase I (Table 1).

Eubacterial Family A DNA polymerases (with proofreading activity)

E. coli DNA pol I (Klenow fragment), *Thermotoga maritima* (UITma fragment)

Family D DNA polymerases (unrelated to Families A, B, C)

Archaeal DNA polymerase II (DP1/DP2) e.g., as described in Cann et al (1998) PNAS 95:14250-5.

Table 1. Accession Information for Cloned Family B Polymerases

Vent *Thermococcus litoralis*

ACCESSION AAA72101

PID g348689

5 VERSION AAA72101.1 GI:348689

DBSOURCE locus THCVDPE accession M74198.1

THEST THERMOCOCCUS SP. (STRAIN TY)

ACCESSION O33845

PID g3913524

10 VERSION O33845 GI:3913524

DBSOURCE swissprot: locus DPOL_THEST, accession O33845

Pab Pyrococcus abyssi

ACCESSION P77916

PID g3913529

15 VERSION P77916 GI:3913529

DBSOURCE swissprot: locus DPOL_PYRAB, accession P77916

PYRHO Pyrococcus horikoshii

ACCESSION O59610

PID g3913526

20 VERSION O59610 GI:3913526

DBSOURCE swissprot: locus DPOL_PYRHO, accession O59610

PYRSE PYROCOCCUS SP. (STRAIN GE23)

ACCESSION P77932

PID g3913530

5 VERSION P77932 GI:3913530

DBSOURCE swissprot: locus DPOL_PYRSE, accession P77932

DeepVent Pyrococcus sp.

ACCESSION AAA67131

PID g436495

10 VERSION AAA67131.1 GI:436495

DBSOURCE locus PSU00707 accession U00707.1

Pfu Pyrococcus furiosus

ACCESSION P80061

PID g399403

15 VERSION P80061 GI:399403

DBSOURCE swissprot: locus DPOL_PYRFU, accession P80061

JDF-3 Thermococcus sp.

Unpublished

Baross gi|2097756|pat|US|5602011|12 Sequence 12 from patent US 5602011

20 9degN THERMOCOCCUS SP. (STRAIN 9ON-7).

ACCESSION Q56366

PID g3913540

VERSION Q56366 GI:3913540

DBSOURCE swissprot: locus DPOL_THES9, accession Q56366

5 KOD *Pyrococcus* sp.

ACCESSION BAA06142

PID g1620911

VERSION BAA06142.1 GI:1620911

DBSOURCE locus PYWKODPOL accession D29671.1

10 Tgo *Thermococcus gorgonarius*.

ACCESSION 4699806

PID g4699806

VERSION GI:4699806

DBSOURCE pdb: chain 65, release Feb 23, 1999

15 THEFM *Thermococcus fumicolans*

ACCESSION P74918

PID g3913528

VERSION P74918 GI:3913528

DBSOURCE swissprot: locus DPOL_THEFM, accession P74918

20 METTH *Methanobacterium thermoautotrophicum*

ACCESSION O27276

PID g3913522

VERSION O27276 GI:3913522

DBSOURCE swissprot: locus DPOL_METTH, accession O27276

5 Metja Methanococcus jannaschii

ACCESSION Q58295

PID g3915679

VERSION Q58295 GI:3915679

DBSOURCE swissprot: locus DPOL_METJA, accession Q58295

10 POC Pyrodictium occultum

ACCESSION B56277

PID g1363344

VERSION B56277 GI:1363344

DBSOURCE pir: locus B56277

15 ApeI Aeropyrum pernix

ACCESSION BAA81109

PID g5105797

VERSION BAA81109.1 GI:5105797

DBSOURCE locus AP000063 accession AP000063.1

20 ARCFU Archaeoglobus fulgidus

ACCESSION O29753

PID g3122019

VERSION O29753 GI:3122019

DBSOURCE swissprot: locus DPOL_ARCFU, accession O29753

5 Desulfurococcus sp. Tok.

ACCESSION 6435708

PID g64357089

VERSION GT:6435708

DBSOURCE pdb. chain 65, release Jun 2, 1999

10 Enzymes possessing 3'-5' exonuclease activity for use in the present compositions and methods may be isolated from natural sources or produced through recombinant DNA techniques. Preferably, the enzyme comprising 3'-5' exonuclease activity is a DNA polymerase.

15 A DNA polymerase comprising 3'-5' exonuclease activity (referred as exo^+) is capable of proofreading the incorporated nucleotides produced by its own polymerization activity. Among other applications, exo^+ DNA polymerases are particularly suited for cloning of PCR products, characterization of polynucleotide sequences. Useful exo^+ DNA polymerases include, but are not limited to, Pwo DNA polymerase; Vent DNA polymerases; Deep Vent DNA polymerase; 9°Nm DNA polymerase; UITma DNA polymerase; Tli DNA polymerase; Pfu DNA polymerase; JDF-3 DNA polymerase; Tgo DNA polymerase; KOD DNA polymerase; and PGB-D DNA
20 polymerase.

In preferred embodiments of the subject invention, an exo^+ DNA polymerase with reduced DNA polymerization activity is used as the second enzyme.

Preparing Exo^+ DNA Polymerase With Reduced DNA Polymerization Activity

25 The cloned wild-type Exo^+ DNA polymerase may be modified to generate forms exhibiting reduced polymerization activity by a number of methods. These include the methods described below and other methods known in the art. Any exo^+ DNA polymerase can be used to prepare for the exo^+ DNA polymerase with reduced DNA polymerization activity in the invention.

5 A. Genetic Modifications - Mutagenesis

The preferred method of preparing a DNA polymerase with reduced polymerization activity is by genetic modification (e.g., by modifying the DNA sequence of a wild-type DNA polymerase). Within the sequence of an exo^+ DNA polymerase, the preferred sequence for genetic modification is the DNA sequence encoding the polymerization domain. Polymerization and exonuclease domains (i.e., their crystal structures) of many DNA polymerases are known in the art (for examples, see Rodriguez et al., 2000, J. Mol. Biol. 299:447-62; Zhao et al., 1999, Structure Fold Des. 7:1189-99; Baker et al., 1998, Proc Natl Acad Sci U S A. 95:3507-12; Kiefer et al., 1997, Structure 5:95-108; Kim et al., 1995, Nature, 376:612-6; Kong et al., 1993, J Biol Chem. 268:1965-75).

General structure features of DNA polymerization domain is known in the art. For example, Blanco et al. (1991, Gene, 100:27-38) discloses that significant amino acid (aa) sequence similarity has been found in the C-terminal portion of 27 DNA-dependent DNA polymerases belonging to the two main superfamilies: (i) Escherichia coli DNA polymerase I (PolI)-like prokaryotic DNA polymerases, and (ii) DNA polymerase alpha-like prokaryotic and eukaryotic (viral and cellular) DNA polymerases. The six most conserved C-terminal regions, spanning approximately 340 amino acids, are located in the same linear arrangement and contain highly conserved motifs and critical residues involved in the polymerization function.

According to the three-dimensional model of PolIk (Klenow fragment), these six conserved regions are located in the proposed polymerization domain, forming the metal and dNTP binding sites and the cleft for holding the DNA template. Site-directed mutagenesis studies support these structural predictions.

The 3'-5' exonuclease active site of E. coli DNA polymerase I is predicted to be conserved for both prokaryotic and eukaryotic DNA polymerases based on amino acid sequence homology (Bernad et al., 1989, Cell, 59:219-28). Three amino acid regions containing the critical residues in the E. coli DNA polymerase I involved in metal binding, single-stranded DNA binding, and catalysis of the exonuclease reaction are located in the amino-terminal half and in the same linear arrangement in several prokaryotic and eukaryotic DNA polymerases. Site-directed mutagenesis at the predicted exonuclease active site of the phi 29 DNA polymerase,

5 a model enzyme for prokaryotic and eukaryotic alpha-like DNA polymerases, specifically
inactivated the 3'-5' exonuclease activity of the enzyme. These results reflect a high evolutionary
conservation of this catalytic domain.

With the great availability of sequences from DNA polymerases, it has become possible
to delineate a few highly conserved regions for various polymerase types (for review, see for
10 example, Johnson, 1993, *Annu Rev Biochem.* 62:685-713). Delarue et al. reported an approach
for unifying the structure of DNA polymerase (1990, *Protein Eng.*, 3:461-7). The speculative
hypothesis should provide a useful model to direct genetic modifications for preparing DNA
polymerase with reduced polymerization activity.

Preferably, the genetic modification for preparing exo^+ DNA polymerase with reduced
15 polymerization activity does not significantly reduces its 3'-5' exonuclease activity (i.e., the
proof reading activity).

Known DNA polymerase mutants that selectively reduce DNA polymerization activity
can be found in the art, for example, in Blanco et al., 1995 *Methods of Enzymology* 262:283-294
((Bacteriophage ϕ 29); Truniger et al., 1996, *EMBO J.* 15:3430-3441 (Bacteriophage ϕ 29);
20 Abdus Sattar et al., 1996, *Biochemistry* 35:16621-9 (Bacteriophage T4); Tuske et al., 2000, *J.*
Biological Chemistry 275:23759-68 (Klenow fragment); Bohlke et al., 2000, *Nucleic Acid*
Research 28:3910-3917 (*Thermococcus aggregans*); Pisani et al., 1998, *Biochemistry* 37:15005-
15012 (*Sulfolobus solfataricus*); Komori et al., 2000, *Protein Eng* 13:41-7 (*Pyrococcus furiosus*);
Shen et al., 2001 *J. Biological Chemistry* 276:27376-83 (*Pyrococcus horikoshi* Family D).

25 Site-directed mutagenesis of bacteriophage ϕ 29 DNA polymerase leads to the
identification of mutations in the polymerase domain which reduce DNA polymerase activity,
while having minimal effects on 3'-5' exonuclease activity (Blanco, L. and Salas, M. 1995,
Methods of Enzymology 262:283-294). In one embodiment of the invention, one or more
corresponding amino acids in Pfu DNA polymerases are mutated (e.g., by substitutions: D405E,
30 Y410F, T542P, D543G, K593T, Y595S). It is understood that other amino acid side
substitutions at these same sites would also selectively reduce DNA polymerase activity.

5 The ϕ 29 DNA polymerase mutagenesis studies targeted amino acid residues within highly conserved Family B motifs (DXXSLYP, KXXXNSXYG, TXXGR, YXDTDS, and KXY), although other regions of the protein presumably can be mutagenized to selectively decrease DNA polymerase activity. One such region is the partitioning domain, characterized by the YXGG motif (Truniger et al., 1996, EMBO J. 15:3430-3441). This region is located within an accessible loop connecting the 3'-5' exonuclease and polymerase domains. The partitioning domain plays a critical role in coordinating the balance between synthesis and degradation of the DNA chain. Mutations within this region disrupt the equilibrium between polymerization and proofreading, and produce phenotypes favoring either polymerization (reduced proofreading) or proofreading (reduced polymerization).

15 Non-conservative (S,N) substitutions at Y₃₈₇ (equivalent to Y₃₈₅ in Pfu) in the partitioning domain of the archaeal *Thermococcus aggregans* DNA polymerase lead to a significant reduction in DNA polymerase activity and enhanced exonuclease activity, which results in improved enzyme fidelity (used alone in PCR) (Bohlke, K. et al (2000) NAR 28:3910-3917). In contrast, conservative substitutions at Y₃₈₇ (F, W, H) lead to wild-type-like fidelity and enhanced PCR performance, which may be related to improved polymerization. A G389A mutation (equivalent to Pfu G387) in *Thermococcus aggregans* DNA polymerase lead to reduced DNA polymerase activity (10% wt), increased exonuclease activity (236% wt), and loss of product synthesis in PCR (Bohlke, K. et al (2000) NAR 28:3910-3917). Analogous mutations have been investigated in bacteriophage ϕ 29 DNA polymerase (Truniger, V., et al (1996) EMBO J. 15:3430-3441) and in the archaeal *Sulfolobus solfataricus* (Sso) DNA polymerase (Pisani, F.M., DeFelice, M., and Rossi, M. (1998) *Biochemistry* 37:15005-15012), where a G→A mutation either decreases (pol/exo = 0.6 for Sso) or increases (pol/exo = 91 for ϕ 29) DNA polymerase activity relative to exonuclease activity.

25 In one embodiment of the invention, Pfu DNA polymerase was mutated within the partitioning domain at amino acids 384-389 (SYTGGF) to obtain a Pfu DNA polymerase with reduced polymerization activity. It is understood that other amino acid side substitutions within the partitioning domain, e.g., at positions Y385, G387, G388, could also selectively reduce DNA polymerase activity while having minimal effects on exonuclease activity.

5 In another embodiment, two or mutations are combined (e.g., by introducing additional site-directed mutations into a mutant Pfu DNA polymerase) to effectively eliminate DNA polymerase activity, while retaining high levels of proofreading activity.

U.S. Patent Nos. 5,691,142, 5,614,402 and 5,541,311 disclose methods of deriving 5'-3' nucleases from thermostable DNA polymerases for the detection of target polynucleotide molecules (hereby incorporated by reference). These methods can be applied to the subject invention for preparing DNA polymerase comprising 3'-5' exonuclease activity with a reduced polymerization activity. Other techniques for genetic modification are well known in the art (see for example, Ausubel et. al. Short Protocols in Molecular Biology (1995) 3rd Ed. John Wiley & Sons, Inc.).

15 Modification to the primary structure of a wild type enzyme by deletion, addition, or alteration of the amino acids incorporated into the sequence during translation can be made without destroying the high temperature DNA polymerase activity of the protein. Such substitutions or other alterations result in proteins useful in the methods of the present invention. The availability of DNA encoding these sequences provides the opportunity to modify the codon sequence to generate mutant enzymes having reduced polymerization activity. A few methods for altering DNA sequences are provided below, any other method known in the art may also be used.

25 There are a number of site-directed mutagenesis methods known in the art which allow one to mutate a particular site or region in a straightforward manner, based on the sequences of the polymerization domain of a DNA polymerase. There are a number of kits available commercially for the performance of site-directed mutagenesis, including both conventional and PCR-based methods. Examples include the EXSITE™ PCR-Based Site-directed Mutagenesis Kit available from Stratagene (Catalog No. 200502) and the QUIKCHANGE™ Site-directed mutagenesis Kit from Stratagene (Catalog No. 200518), and the CHAMELEON® double-stranded Site-directed mutagenesis kit, also from Stratagene (Catalog No. 200509).

Older methods of site-directed mutagenesis known in the art relied upon sub-cloning of the sequence to be mutated into a vector, such as an M13 bacteriophage vector, that allows the isolation of single-stranded DNA template. In these methods one anneals a mutagenic primer

5 (i.e., a primer capable of annealing to the site to be mutated but bearing one or mismatched nucleotides at the site to be mutated) to the single-stranded template and then polymerizes the complement of the template starting from the 3' end of the mutagenic primer. The resulting duplexes are then transformed into host bacteria and plaques are screened for the desired mutation.

10 More recently, site-directed mutagenesis has employed PCR methodologies, which have the advantage of not requiring a single-stranded template. In addition, methods have been developed that do not require sub-cloning. Several issues must be considered when PCR-based site-directed mutagenesis is performed. First, in these methods it is desirable to reduce the number of PCR cycles to prevent expansion of undesired mutations introduced by the
15 polymerase. Second, a selection must be employed in order to reduce the number of non-mutated parental molecules persisting in the reaction. Third, an extended-length PCR method is preferred in order to allow the use of a single PCR primer set. And fourth, because of the non-template-dependent terminal extension activity of some thermostable polymerases it is often necessary to incorporate an end-polishing step into the procedure prior to blunt-end ligation of
20 the PCR-generated mutant product.

The protocol described below accommodates these considerations through the following steps. First, the template concentration used is approximately 1000-fold higher than that used in conventional PCR reactions, allowing a reduction in the number of cycles from 25-30 down to 5-
10 without dramatically reducing product yield. Second, the restriction endonuclease DpnI
25 (recognition target sequence: 5-Gm6ATC-3, where the A residue is methylated) is used to select against parental DNA, since most common strains of E. coli Dam methylate their DNA at the sequence 5-GATC-3. Third, Taq Extender is used in the PCR mix in order to increase the proportion of long (i.e., full plasmid length) PCR products. Finally, Pfu DNA polymerase is used to polish the ends of the PCR product prior to intramolecular ligation using T4 DNA ligase.

30 A non-limiting example for the method is described in detail as follows:

Plasmid template DNA (approximately 0.5 pmole) is added to a PCR cocktail containing: 1x mutagenesis buffer (20 mM Tris HCl, pH 7.5; 8 mM MgCl₂; 40 µg/ml BSA); 12-20 pmole of each primer (one of skill in the art may design a mutagenic primer as necessary, giving

5 consideration to those factors such as base composition, primer length and intended buffer salt concentrations that affect the annealing characteristics of oligonucleotide primers; one primer must contain the desired mutation, and one (the same or the other) must contain a 5' phosphate to facilitate later ligation), 250 μ M each dNTP, 2.5 U Taq DNA polymerase, and 2.5 U of Taq Extender (Available from Stratagene; See Nielson et al. (1994) Strategies 7: 27, and U.S. Patent
10 No. 5,556,772). Primers can be prepared using the triester method of Matteucci et al., 1981, J. Am. Chem. Soc. 103:3185-3191, incorporated herein by reference. Alternatively automated synthesis may be preferred, for example, on a Biosearch 8700 DNA Synthesizer using cyanoethyl phosphoramidite chemistry.

15 The PCR cycling is performed as follows: 1 cycle of 4 min at 94°C, 2 min at 50°C and 2 min at 72°C; followed by 5-10 cycles of 1 min at 94°C, 2 min at 54°C and 1 min at 72°C. The parental template DNA and the linear, PCR-generated DNA incorporating the mutagenic primer are treated with DpnI (10 U) and Pfu DNA polymerase (2.5U). This results in the DpnI digestion of the in vivo methylated parental template and hybrid DNA and the removal, by Pfu DNA polymerase, of the non-template-directed Taq DNA polymerase-extended base(s) on the
20 linear PCR product. The reaction is incubated at 37°C for 30 min and then transferred to 72°C for an additional 30 min. Mutagenesis buffer (115 μ l of 1x) containing 0.5 mM ATP is added to the DpnI-digested, Pfu DNA polymerase-polished PCR products. The solution is mixed and 10 μ l are removed to a new microfuge tube and T4 DNA ligase (2-4 U) is added. The ligation is incubated for greater than 60 min at 37°C. Finally, the treated solution is transformed into
25 competent E. coli according to standard methods.

Methods of random mutagenesis which will result in a panel of mutants bearing one or more randomly-situated mutations exist in the art. Such a panel of mutants may then be screened for those exhibiting reduced polymerization relative to the wild-type polymerase (e.g., by measuring the incorporation of 10nmoles of dNTPs into polymeric form in 30 minutes at the
30 optimal temperature for a given DNA polymerase). An example of a method for random mutagenesis is the so-called "error-prone PCR method". As the name implies, the method amplifies a given sequence under conditions in which the DNA polymerase does not support high fidelity incorporation. The conditions encouraging error-prone incorporation for different DNA polymerases vary, however one skilled in the art may determine such conditions for a

5 given enzyme. A key variable for many DNA polymerases in the fidelity of amplification is, for example, the type and concentration of divalent metal ion in the buffer and the inherited fidelity of the PCR enzyme. The use of manganese ion and/or variation of the magnesium or manganese ion concentration may therefore be applied to influence the error rate of the polymerase.

10 In a preferred embodiment, the second enzyme with reduced polymerization activity is derived from Pfu DNA polymerase.

The DNA coding sequence of a wild-type Pfu DNA polymerase can be found in the art, for example, from Genbank (accession No. U84155). A detailed description of the structure and function of Pfu DNA polymerase can be found, among other places in U.S. Patent Nos. 5,948,663; 5,866,395; 5,545,552; 5,556,772, all of which are hereby incorporated by reference. A not-limiting detailed procedure for preparing Pfu DNA polymerase with reduced polymerization activity is provided in Example 1.

15 A person of average skill in the art having the benefit of this disclosure will recognize that polymerases with reduced polymerization activity derived from other exo^+ DNA polymerases including Vent DNA polymerase, JDF-3 DNA polymerase, Tgo DNA polymerase and the like may be suitably used in the subject compositions.

20 The first or the second enzyme of the subject composition may comprise DNA polymerases that have not yet been isolated. Assays for both DNA polymerization activity and 3'-5' exonuclease activity can be found in the subject description and in DNA Replication 2nd Ed., Kornberg and Baker, supra; Enzymes, Dixon and Webb, Supra, as well as other publications available to the person of ordinary skill in the art.

In preferred embodiments of the invention, mutant Pfu DNA polymerase comprises one or more mutations at amino acid positions selected from the group consisting of: D405, Y410, T542, D543, K593, Y595, Y385, G387, and G388.

30 More preferably, the mutant Pfu DNA polymerase comprises one or more mutations selected from the group consisting of: D405E, Y410F, T542P, D543G, K593T, Y595S, Y385Q, Y385S, Y385N, Y385L, Y385H, G387S, G387P, and G388P.

The invention encompasses compositions and methods in which a mutant of a related archaeal DNA polymerase is with reduced (e.g., deficient in) polymerase activity, while retaining proofreading activity. Such mutations may be within the partitioning domain or the polymerase domain of the DNA polymerases. Table 2 (A and B) provides an unlimited example of such mutations in various DNA polymerases. A mutant DNA polymerase of the invention may comprise a single mutation as indicted in Table 2, or a combination of any two or more mutations.

Table 2A Partitioning Domain Mutations in Various DNA Polymerases

Enzyme	Domain (bp)	Domain sequence	Predicted Mutations for Reducing DNA Polymerase Activity*	Preferred mutation
Pfu	384-389	S Y T G G F	Y385, G387, G388 (Y385N, Y385L, Y385H, Y385Q, Y385S; G387S, G387P; G388A, G388P)	G387P
Tgo	383-388	S Y A G G Y	Y384, G386, G387 (Y384N, Y384L, Y384H, Y384Q, Y384S; G386S, G386P; G387A, G387P)	G386P
KOD	383-388	S Y E G G Y	Y384, G386, G387 (Y384N, Y384L, Y384H, Y384Q, Y384S; G386S, G386P; G387A, G387P)	G386P
Vent	386-391	T Y L G G Y	Y387, G389, G390 (Y387N, Y387L, Y387H, Y387Q, Y387S; G389S, G389P; G390A, G390P)	G389P

DeepVent	384-389	S Y A G G Y	Y385, G387, G388 (Y385N, Y385L, Y385H, Y385Q, Y385S; G387S, G387P; G388A, G388P)	G387P
----------	---------	-------------	---	-------

5

Table 2B Polymerase Domain Mutations in Various DNA Polymerases

Enzyme	Domain (bp)	Domain sequence	Predicted Mutations for Reducing DNA Polymerase Activity#
Pfu	DXXSLYP 405-411	DFRALYP	D405 (D405E)
Tgo	404-410	DFRSLYP	D404 (D404E)
KOD	404-410	DFRSLYP	D404 (D404E)
Vent	407-413	DFRSLYP	D407 (D404E)
Deep Vent	405-411	DFRSLYP	D405 (D404E)
Pfu	YXDTDS 539-544	YIDTDG	T542, D543 (T542P; D543G)
Tgo	538-543	YADTDG	T541, D542 (T541P; D542G)
KOD	538-543	YSDDTG	T541, D542 (T541P; D542G)
Vent	541-546	YADTDG	T544, D545 (T544P; D545G)
Deep Vent	539-544	YIDTDG	T542, D543 (T542P; D543G)
Pfu	KXY 593-595	KRY	K593 (K593T)
Tgo	592-594	KKY	K592 (K592T)
KOD	592-594	KKY	K592 (K592T)

Vent	595-597	KRY	K595 (K595T)
Deep Vent	593-595	KKY	K593 (K593T)

5 #alternative side chain substitutions at key positions are also expected to reduce polymerase activity

B. Methods of Evaluating Mutants for Reduced Polymerization

Random or site-directed mutants generated as known in the art or as described herein and expressed in bacteria may be screened for reduced polymerization by several different assays.

10 Embodiments for the expression of mutant and wild type enzymes is described herein below in section C. In one method, exo^+ DNA polymerase proteins expressed in lytic lambda phage plaques generated by infection of host bacteria with expression vectors based on, for example, Lambda ZapII[®], are transferred to a membrane support. The immobilized proteins are then assayed for polymerase activity on the membrane by immersing the membranes in a buffer
15 containing a DNA template and the unconventional nucleotides to be monitored for incorporation.

Mutant polymerase libraries may be screened using a variation of the technique used by Sagner et al (Sagner, G., Ruger, R., and Kessler, C. (1991) Gene 97:119-123). For this approach, lambda phage clones are plated at a density of 10-20 plaques per square centimeter. Proteins
20 present in the plaques are transferred to filters and moistened with polymerase screening buffer (50mM Tris (pH 8.0), 7mM MgCl₂, 3mM β -ME). The filters are kept between layers of plastic wrap and glass while the host cell proteins are heat-inactivated by incubation at 65°C for 30 minutes. The heat-treated filters are then transferred to fresh plastic wrap and approximately 351 of polymerase assay cocktail are added for every square centimeter of filter. The assay cocktail
25 consists of 1X cloned Pfu (cPfu) magnesium free buffer (1X buffer is 20 mM Tris-HCl (pH 8.8), 10 mM KCl, 10 mM (NH₄)₂SO₄, 100 μ g/ml bovine serum albumin (BSA), and 0.1% Triton X-100; Pfu Magnesium-free buffer may be obtained from Stratagene (Catalog No. 200534)), 125 ng/ml activated calf thymus or salmon sperm DNA, 1.29 μ Ci/ml α -³³P ddNTP. The filters are placed between plastic wrap and a glass plate and then incubated at 65°C for one hour, and then
30 at 70°C for one hour and fifteen minutes. Filters are then washed three times in 2X SSC for five minutes per wash before rinsing twice in 100% ethanol and vacuum drying. Filters are then

5 exposed to X-ray film (approximately 16 hours), and plaques that incorporate label are identified by aligning the filters with the original plate bearing the phage clones. Plaques identified in this way are re-plated at more dilute concentrations and assayed under similar conditions to allow the isolation of purified plaques.

10 In assays such as the one described above, the signal generated by the label is a direct measure of the polymerization activity of the polymerase. A plaque comprising a mutant DNA polymerase with reduced DNA polymerization activity compared to that of the wild-type enzyme can be selected.

15 Incorporation of nucleotides may also be measured in extension reactions by adding, for example, 1 µl of appropriately diluted bacterial extract (i.e., heat-treated and clarified extract of bacterial cells expressing a cloned polymerase or mutated cloned polymerase) to 10 µl of each nucleotide cocktail, followed by incubation at the optimal temperature for 30 minutes (e.g., 73°C for Pfu DNA polymerase), for example, as described in Hogrefe et al., 2001, Methods in Enzymology, 343:91-116. Extension reactions are quenched on ice, and then 5µl aliquots are spotted immediately onto DE81 ion-exchange filters (2.3cm; Whatman #3658323).
20 Unincorporated label is removed by 6 washes with 2xSCC (0.3M NaCl, 30mM sodium citrate, pH 7.0), followed by a brief wash with 100% ethanol. Incorporated radioactivity is then measured by scintillation counting. Reactions that lack enzyme are also set up along with sample incubations to determine “total cpms” (omit filter wash steps) and “minimum cpms”(wash filters as above). Cpms bound is proportional to the amount of polymerase activity present per volume
25 of bacterial extract.

A Non-limiting method for determining polymerization activity of a DNA polymerase mutant relative to wild type (wt) is provided as follows. Relative percent radioactivity incorporated which indicates the relative polymerization activity of a DNA polymerase mutant can be determined as:

$$\frac{(\text{corrected cpms for mutant DNA polymerase}) \times (\text{ng wt DNA polymerase})}{(\text{corrected cpms for wt DNA polymerase}) \times (\text{ng mutant DNA polymerase})}$$

To more precisely quantify % activity, one should convert cpms incorporated into units of DNA polymerase activity. One unit of polymerase activity is defined as the amount of enzyme that catalyzes the incorporation of 10nmols of total dNTP into polymeric form (e.g., binds to DE-81 paper) in 30 minutes at optimal temperature. Units of DNA polymerase activity can be calculated using the following equation:

$$\frac{(\text{corrected sample cpms})}{\text{total cpms}} \times \frac{(8\text{nmols dNTPs})}{\text{reaction}} \times \frac{(1 \text{ unit})}{(10\text{nmols dNTPs incorporated})}$$

Polymerase specific activity (U/mg) can be extrapolated from the slope of the linear portion of units versus enzyme amount plots. Protein concentrations can be determined relative to a BSA standard (Pierce) in a colorimetric assay (e.g. Pierce's Coomassie Plus Protein Assay). Alternatively, when protein amounts are limiting (or for preparations of limited purity), relative protein concentrations can be verified by SDS-PAGE analysis. Several aliquots of each DNA polymerase preparation, ranging from 1-20 ng of total protein, are subject to SDS-PAGE electrophoresis and the intensity of silver- and/or Sypro orange (Molecular Probes)-stained bands are compared to standards. Finally, % activity can be determined as:

$$\frac{\text{specific polymerase activity (U/mg) of mutant DNA polymerase}}{\text{specific polymerase activity (U/mg) of wt DNA polymerase}}$$

It is preferred that the polymerases with reduced polymerization activity of the present invention maintain their proofreading activities (i.e., 3'-5' exonuclease activities). The mutant DNA polymerases with reduced DNA polymerization activities, therefore, are also assayed for 3'-5- exonuclease activities.

Suitable exonuclease activity assays include one described in Hogrefe et al (supra, and as described in Example 3). Another assay employs double-stranded λ DNA, which has been uniformly labeled with ^3H S-adenosyl methionine (NEN #NET-155) and *Sss* I methylase (NEB), and then restriction digested with *Pal* I (Kong et al., 1993, J. Biol. Chem. 268:1965). Using double-stranded labeled DNA templates, one can determine specificity by measuring whether cpms decrease (3'-5' exonuclease) with the addition of dNTPs (10-100 μM). A typical exonuclease reaction cocktail consists of 1x reaction buffer and 20 $\mu\text{g/ml}$ ^3H -labeled digested

5 double-stranded λ DNA ($\sim 10^6$ cpms/ml), prepared as described (Kong et al., supra).
 Exonuclease activity can be measured in the appropriate PCR buffer or in a universal assay
 buffer such as 70mM Tris HCl (pH 8.8), 2mM MgCl₂, 0.1% Triton-X, and 100 μ g/ml BSA.

Percent exonuclease activity can be determined as: (corrected cpms for
 mutants)/(corrected cpms for wt DNA polymerase). To more precisely quantify % activity,
 10 cpms released can be converted into units of exonuclease activity. One unit of exonuclease
 activity is defined as the amount of enzyme that catalyzes the acid-solubilization of 10nmoles of
 total dNMPs in 30 minutes at a defined temperature. To determine units, background (average
 “minimum cpms” value) is first subtracted from the average sample cpms. Nmoles dNMPs
 released is calculated using the following equation:

$$\frac{(\text{corrected sample cpms})}{\text{total cpms}} \times \frac{(920\text{ng DNA})}{\text{reaction}} \times \frac{(1\text{n mole dNMP})}{(330\text{ng dNMP})}$$

Units of exonuclease activity (in 30 minutes) can then be determined as:

$$\frac{(\text{nmoles dNMPs released per hr})}{2} \times \frac{(1 \text{ unit})}{(10\text{nmoles dNMPs released})}$$

Exonuclease specific activity (U/mg) can be extrapolated from the slope of the linear portion of
 units versus enzyme amount plots. Finally, % activity can be determined as:

$$\frac{\text{specific exonuclease activity (U/mg) of mutant DNA polymerase}}{\text{specific exonuclease activity (U/mg) of wt DNA polymerase}}$$

In addition to the substrate described above, exonuclease activity can be also be
 quantified using [³H]-*E. coli* genomic DNA (NEN #NET561; 5.8 μ Ci/ μ g), a commercially-
 available substrate. A typical exonuclease reaction cocktail consists of 0.4 μ g/ml ³H-labeled *E.*
 30 *coli* genomic DNA in 1x reaction buffer. Assays are performed as described above.

5 Genes for desired mutant DNA polymerases generated by mutagenesis may be sequenced
to identify the sites and number of mutations. For those mutants comprising more than one
mutation, the effect of a given mutation may be evaluated by introduction of the identified
mutation to the wild-type gene by site-directed mutagenesis in isolation from the other mutations
borne by the particular mutant. Screening assays of the single mutant thus produced will then
10 allow the determination of the effect of that mutation alone.

In one embodiment, the Pfu mutant is G387P, which reduces the error rate of wild type
Pfu DNA polymerase by 3-fold in Pfu blend and the error rate of Taq by 5- to 8-fold in Taq
blend when added at 0.6-3.6µg/50µl reaction. Pfu G387P exhibited 0.4% DNA polymerase
activity and 57% exonuclease activity (i.e., relative to wild type Pfu) in a preliminary screen of
15 partially purified (~50% purity) His-tagged proteins, eluted from nickel columns (Table 1).
After column chromatography (~95% purity), the Pfu G387P mutant was found to be devoid of
detectable DNA polymerase activity (<0.01% activity relative to wild type Pfu) (Table 3).

C. Expression of Wild-type or Mutant enzymes According to the Invention

Methods known in the art may be applied to express and isolate the mutated forms of
DNA polymerase (i.e., the second enzyme) according to the invention. The methods described
20 here can be also applied for the expression of wild-type enzymes useful (e.g., the first enzyme) in
the invention. Many bacterial expression vectors contain sequence elements or combinations of
sequence elements allowing high level inducible expression of the protein encoded by a foreign
sequence. For example, as mentioned above, bacteria expressing an integrated inducible form of
25 the T7 RNA polymerase gene may be transformed with an expression vector bearing a mutated
DNA polymerase gene linked to the T7 promoter. Induction of the T7 RNA polymerase by
addition of an appropriate inducer, for example, isopropyl-β-D-thiogalactopyranoside (IPTG) for
a lac-inducible promoter, induces the high level expression of the mutated gene from the T7
promoter.

30 Appropriate host strains of bacteria may be selected from those available in the art by one
of skill in the art. As a non-limiting example, E. coli strain BL-21 is commonly used for
expression of exogenous proteins since it is protease deficient relative to other strains of E. coli.
BL-21 strains bearing an inducible T7 RNA polymerase gene include WJ56 and ER2566

5 (Gardner & Jack, 1999, supra). For situations in which codon usage for the particular
polymerase gene differs from that normally seen in E. coli genes, there are strains of BL-21 that
are modified to carry tRNA genes encoding tRNAs with rarer anticodons (for example, argU,
ileY, leuW, and proL tRNA genes), allowing high efficiency expression of cloned protein genes,
for example, cloned archaeal enzyme genes (several BL21-CODON PLUSTM cell strains
10 carrying rare-codon tRNAs are available from Stratagene, for example).

There are many methods known to those of skill in the art that are suitable for the
purification of a modified DNA polymerase of the invention. For example, the method of
Lawyer et al. (1993, PCR Meth. & App. 2: 275) is well suited for the isolation of DNA
polymerases expressed in E. coli, as it was designed originally for the isolation of Taq
polymerase. Alternatively, the method of Kong et al. (1993, J. Biol. Chem. 268: 1965,
15 incorporated herein by reference) may be used, which employs a heat denaturation step to
destroy host proteins, and two column purification steps (over DEAE-Sepharose and heparin-
Sepharose columns) to isolate highly active and approximately 80% pure DNA polymerase.
Further, DNA polymerase mutants may be isolated by an ammonium sulfate fractionation,
followed by Q Sepharose and DNA cellulose columns, or by adsorption of contaminants on a
20 HiTrap Q column, followed by gradient elution from a HiTrap heparin column.

In one embodiment, the Pfu mutants are expressed and purified as described in U.S.
Patent No. 5,489,523, thereby incorporated by reference in its entirety.

D. Other Methods For Reducing Polymerization Activity

25 In order to prevent the side effects of having a high DNA polymerization activity in an
amplification reaction, the polymerization activity of the composition of the invention may also
be reduced by physical and/or chemical modification and/or inhibition.

The polymerization activity of the subject composition may be reduced by chemical
and/or physical means. Conditions which preferentially inhibit the polymerization activity of a
30 DNA polymerase is known in the art (for reviews, see Johnson, 1993, supra; Wright, 1996, Acta
Biochim Pol. 43:115-24; Elion, 1982, Am J Med., 73:7-13). The level of polymerization activity
need only be reduced to that level of activity which does not interfere with amplification

5 reactions (e.g., does not significantly affect the exo^+ activity of the composition or the efficiency yield of the amplification reaction).

Concentrations of Mg^{2+} greater than 5 mM inhibit the polymerization activity of the Pfu DNA polymerase. The effect of a given concentration of Mg^{2+} for a given DNA polymerase may be determined by quantitation of the efficiency and specificity of polymerization.

10 The inhibitory effect of other ions, polyamines, denaturants, such as urea, formamide, dimethylsulfoxide, glycerol and non-ionic detergents (Triton X-100 and Tween-20), polynucleotide binding chemicals such as, actinomycin D, ethidium bromide and psoralens, may be tested by their addition to the standard reaction buffers for polynucleotide amplification (e.g., PCR). Those compounds having a preferential inhibitory effect on the polymerization activity
15 but not significantly affecting the 3'-5' exonuclease activity of a DNA polymerase are then used to create reaction conditions under which 3'-5' nuclease activity is retained while polymerization activity is reduced.

Physical means may be used to preferentially inhibit the polymerization activity of a polymerase. For example, the polymerization activity of thermostable polymerases is destroyed
20 by exposure of the polymerase to extreme heat (typically 96°C to 100°C) for extended periods of time (greater than or equal to 20 minutes). While there are minor differences with respect to the specific heat tolerance for each of the enzymes, these are readily determined. The polymerase mixture of the invention or the exo^+ DNA polymerase used as the second enzyme with reduced polymerization activity can be treated with heat for various periods of time and the effect of the
25 heat treatment upon the polymerization and 3'-5' nuclease activities is determined. Conditions reducing DNA polymerase activity but not significantly affecting the 3'-5' exonuclease activity may be used to pretreat the polymerase mixture or the exo^+ DNA polymerase used as second enzyme with reduced polymerization activity in the present invention.

Enzyme Mixture

30 The subject enzyme mixture composition comprises a first enzyme comprising DNA polymerization activity and a second enzyme comprising 3'-5' exonuclease activity with reduced DNA polymerase activity.

5 In one embodiment, the first enzyme is a DNA polymerase with 3'-5' exonuclease activity. The fidelity of the first enzyme for DNA amplification is increased by the use of a second enzyme which also possesses 3'-5' exonuclease activity. A preferred DNA polymerase with 3'-5' exonuclease activity as the first enzyme is a wild type Pfu DNA polymerase.

10 In another embodiment, the first enzyme is a DNA polymerase without 3'-5' exonuclease activity. The fidelity of an amplification reaction is provided by the second enzyme of the subject invention, which possesses 3'-5' exonuclease activity. A preferred DNA polymerase without 3'-5' exonuclease activity as the first enzyme is a Taq DNA polymerase.

15 In yet another embodiment, the first enzyme may be a reverse transcriptase with DNA polymerization activity. The fidelity of the reverse transcriptase in cDNA synthesis is increased by the use of a second enzyme which possesses 3'-5' exonuclease activity.

A. Selection of the first and the second enzyme pair

In the subject method for DNA synthesis, any enzyme comprising DNA polymerization activity may be mixed with a second enzyme comprising 3'-5' exonuclease activity and reduced polymerization activity.

20 When both first and second enzymes in the mixture comprise 3'-5' exonuclease activity, it may be desirable to combine two enzymes with different proofreading activities. By "different proofreading activity", it means that two 3'-5' exonucleases exhibit different proofreading preference for a nucleotide. For example, one 3'-5' exonuclease may proofread a G-T mismatch more efficiently than an A-A mismatch, another exonuclease having a different proofreading preference may proofread an A-A mismatch more efficiently than a G-T mismatch. By using a
25 second enzyme with a different proofreading preference from the first enzyme of the subject composition, one can enhance proofreading of the first enzyme by providing proofreading to mismatches which the first enzyme is not capable of recognizing and excising efficiently.

30 Another factor to consider when selecting the first and the second enzymes of the subject invention is the compatibility of reaction conditions (e.g., pH, buffer composition, temperature requirement, etc.) required by each enzyme.

5 In a preferred embodiment, the subject composition comprises a wild-type Pfu DNA polymerase as the first enzyme and a mutant Pfu DNA polymerase with reduced DNA polymerization activity as the second enzyme. Preferably, the mixture comprises a ratio of 2.5-5U Pfu DNA polymerase plus an amount of a polymerase reduced mutant corresponding to <0.1U DNA polymerase activity and 0.05U to 10U of 3'-5' exonuclease activity (or the amount of exonuclease activity containing within 2.5-500U wild type Pfu). More preferably, the mixture comprises a ratio of 2.5-5U Pfu DNA polymerase plus an amount of a polymerase reduced mutant corresponding to <0.01U DNA polymerase activity and 0.5U to 1.7U of 3'-5' exonuclease activity (or the amount of exonuclease activity contained within 30-90U wild type Pfu)

10 In another preferred embodiment, the subject composition comprises a wild-type Taq DNA polymerase as the first enzyme and a mutant Pfu DNA polymerase with reduced DNA polymerization activity as the second enzyme. Preferably, the enzyme mixture comprises a ration of 2.5U Taq DNA polymerase plus an amount of a polymerase deficient mutant corresponding to <0.00125U DNA polymerase activity and 0.05U to 10U of 3'-5' exonuclease activity (or the amount of exonuclease activity contained within 2.5-500U wild type Pfu). More preferably, the enzyme mixture comprises a ratio of 2.5U Taq DNA polymerase plus an amount of a polymerase deficient mutant corresponding to <0.00125U DNA polymerase activity and 0.5U to ≥ 3 U of 3'-5' exonuclease activity (or the amount of exonuclease activity contained within 30 to ≥ 160 U wild type Pfu).

25 Preferably the mutant Pfu DNA polymerase with reduced DNA polymerization activity comprises one or more mutations at amino acid positions selected from the group consisting of: D405, Y410, T542, D543, K593, Y595, Y385, G387, and G388.

More preferably, the mutant Pfu DNA polymerase comprises one or more mutations selected from the group consisting of: D405E, Y410F, T542P, D543G, K593T, Y595S, Y385Q, Y385S, Y385N, Y385L, Y385H, G387S, G387P, and G388P.

5 B. The Ratio Of Polymerization To Exonuclease Activity In The Enzyme Mixture

In a variety of DNA synthesis and amplification procedures, the compositions of the present invention provide superior synthesis results (e.g., higher fidelity and efficiency), as compared with the synthesis results obtained with a single DNA polymerase or with a mixture comprising two wild type DNA polymerases. When using the subject composition, the ratio of
10 total polymerization activity and total exonuclease activity in the enzyme mixture may be critical for optimal efficiency and fidelity of DNA synthesis.

In the enzyme mixture of the subject invention, when DNA polymerases are used as the first and second enzymes, both enzymes may contribute to the polymerization and/or 3'-5' exonuclease activity. When an enzyme other than a conventional DNA polymerase is used as
15 the first enzyme (e.g., a reverse transcriptase), both enzymes may contribute to DNA polymerization activity, but only the second enzyme contribute to the 3'-5' exonuclease activity. When an enzyme other than a DNA polymerase is used as the second enzyme (e.g., E. coli exonuclease I), both enzymes may contribute to the 3'-5' exonuclease activity, but only the first
20 enzyme contribute to the polymerization activity of the enzyme mixture.

The ratio of the first and the second enzyme in the subject composition may be varied with respect to one another. The ratio of the DNA polymerization activity to 3'-5' exonuclease activity present in the subject composition employed in a given synthesis procedure may be readily optimized by performing a series of simple experiments in which the ratio of the DNA
25 polymerization activity to the exonuclease activity in the reaction mixture are systematically varied with respect to one another and the synthesis results compared.

3'-5' exonuclease activity has been shown to degrade unannealed primers. The degraded primers would not be available in subsequent rounds of DNA amplification and would therefore effect the efficiency of the PCR reaction. In applications requiring very high product yield, it may therefore be desirable to have a low concentration of the exonuclease activity relative to the
30 DNA polymerization activity to decrease this effect and to increase the product yield. However, when fidelity is more important than yield, it may be desirable to have a high concentration of the exonuclease activity relative to the DNA polymerization activity to increase the accuracy of

5 the synthesis or amplification so long as the level of polymerization activity does not significantly inhibit the efficiency of the amplification.

In a preferred embodiment, the ratio of the DNA polymerase activity and the exonuclease activity in the enzyme mixture is about (2.5-5U of DNA polymerization activity)/(0.05-10U of 3'-5' exonuclease activity).

10 Applications of The Subject Invention

In one aspect, the invention provides a method for DNA synthesis using the compositions of the subject invention. The subject compositions may be used in various methods of polynucleotide synthesis in essentially the same manner as the DNA polymerase or other synthetic enzyme present in the subject composition. Typically, synthesis of a polynucleotide requires a synthesis primer, a synthesis template, polynucleotide precursors for incorporation into the newly synthesized polynucleotide, (e.g. dATP, dCTP, dGTP, dTTP), and the like. Detailed methods for carrying out polynucleotide synthesis are well known to the person of ordinary skill in the art and can be found, for example, in Molecular Cloning second edition, Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

20 A. Application In Amplification Reactions

"Polymerase chain reaction" or "PCR" refers to an in vitro method for amplifying a specific polynucleotide template sequence. The technique of PCR is described in numerous publications, including, PCR: A Practical Approach, M. J. McPherson, et al., IRL Press (1991), PCR Protocols: A Guide to Methods and Applications, by Innis, et al., Academic Press (1990), and PCR Technology: Principals and Applications for DNA Amplification, H. A. Erlich, Stockton Press (1989). PCR is also described in many U.S. Patents, including U.S. Patent Nos. 4,683,195; 4,683,202; 4,800,159; 4,965,188; 4,889,818; 5,075,216; 5,079,352; 5,104,792; 5,023,171; 5,091,310; and 5,066,584, each of which is herein incorporated by reference.

For ease of understanding the advantages provided by the present invention, a summary of PCR is provided. The PCR reaction involves a repetitive series of temperature cycles and is typically performed in a volume of 50-100 µl. The reaction mix comprises dNTPs (each of the four deoxynucleotides dATP, dCTP, dGTP, and dTTP), primers, buffers, DNA polymerase, and

5 polynucleotide template. PCR requires two primers that hybridize with the double-stranded target polynucleotide sequence to be amplified. In PCR, this double-stranded target sequence is denatured and one primer is annealed to each strand of the denatured target. The primers anneal to the target polynucleotide at sites removed from one another and in orientations such that the extension product of one primer, when separated from its complement, can hybridize to the other
10 primer. Once a given primer hybridizes to the target sequence, the primer is extended by the action of a DNA polymerase. The extension product is then denatured from the target sequence, and the process is repeated.

In successive cycles of this process, the extension products produced in earlier cycles serve as templates for DNA synthesis. Beginning in the second cycle, the product of
15 amplification begins to accumulate at a logarithmic rate. The amplification product is a discrete double-stranded DNA molecule comprising: a first strand which contains the sequence of the first primer, eventually followed by the sequence complementary to the second primer, and a second strand which is complementary to the first strand.

Due to the enormous amplification possible with the PCR process, small levels of DNA carryover from samples with high DNA levels, positive control templates or from previous
20 amplifications can result in PCR product, even in the absence of purposefully added template DNA. If possible, all reaction mixes are set up in an area separate from PCR product analysis and sample preparation. The use of dedicated or disposable vessels, solutions, and pipettes (preferably positive displacement pipettes) for RNA/DNA preparation, reaction mixing, and
25 sample analysis will minimize cross contamination. See also Higuchi and Kwok, 1989, *Nature*, 339:237-238 and Kwok, and Orrego, in: Innis et al. eds., 1990, PCR Protocols: A Guide to Methods and Applications, Academic Press, Inc., San Diego, Calif., which are incorporated herein by reference.

1. Thermostable Enzymes

30 For PCR amplifications, the enzymes used in the invention are preferably thermostable. As used herein, "thermostable" refers to an enzyme which is stable to heat, is heat resistant, and functions at high temperatures, e.g., 50 to 90°C. The thermostable enzyme according to the present invention must satisfy a single criterion to be effective for the amplification reaction, i.e.,

5 the enzyme must not become irreversibly denatured (inactivated) when subjected to the elevated temperatures for the time necessary to effect denaturation of double-stranded polynucleotides. By “irreversible denaturation” as used in this connection, is meant a process bringing a permanent and complete loss of enzymatic activity. The heating conditions necessary for denaturation will depend, e.g., on the buffer salt concentration and the length and nucleotide
10 composition of the polynucleotides being denatured, but typically range from 85°C, for shorter polynucleotides, to 105°C for a time depending mainly on the temperature and the polynucleotide length, typically from 0.25 minutes for shorter polynucleotides, to 4.0 minutes for longer pieces of DNA. Higher temperatures may be tolerated as the buffer salt concentration and/or GC composition of the polynucleotide is increased. Preferably, the enzyme will not
15 become irreversibly denatured at 90 to 100°C. An enzyme that does not become irreversibly denatured, according to the invention, retains at least 10%, or at least 25%, or at least 50% or more function or activity during the amplification reaction.

2. PCR Reaction Mixture

In addition to the subject enzyme mixture, one of average skill in the art may also employ
20 other PCR parameters to increase the fidelity of synthesis/amplification reaction. It has been reported PCR fidelity may be affected by factors such as changes in dNTP concentration, pH, units of enzyme used per reaction, and the ratio of Mg^{2+} to dNTPs present in the reaction (Mattila et al., 1991, supra).

Mg^{2+} concentration affects the annealing of the oligonucleotide primers to the template
25 DNA by stabilizing the primer-template interaction, it also stabilizes the replication complex of polymerase with template-primer. It can therefore also increase non-specific annealing and produced undesirable PCR products (gives multiple bands in gel). When non-specific amplification occurs, Mg^{2+} may need to be lowered or EDTA can be added to chelate Mg^{2+} to increase the accuracy and specificity of the amplification.

30 Other divalent cations such as Mn^{2+} , or Co^{2+} can also affect DNA polymerization. Suitable cations for each DNA polymerase are known in the art (e.g., in DNA Replication 2nd edition, supra). Divalent cation is supplied in the form of a salt such $MgCl_2$, $Mg(OAc)_2$, $MgSO_4$, $MnCl_2$, $Mn(OAc)_2$, or $MnSO_4$. Usable cation concentrations in a Tris-HCl buffer are for $MnCl_2$

5 from 0.5 to 7 mM, preferably, between 0.5 and 2 mM, and for MgCl_2 from 0.5 to 10 mM. Usable cation concentrations in a Bicine/KOAc buffer are from 1 to 20 mM for $\text{Mn}(\text{OAc})_2$, preferably between 2 and 5 mM.

10 Monovalent cation required by DNA polymerase may be supplied by the potassium, sodium, ammonium, or lithium salts of either chloride or acetate. For KCl, the concentration is between 1 and 200 mM, preferably the concentration is between 5 and 100 mM, although the optimum concentration may vary depending on the polymerase used in the reaction.

15 Deoxyribonucleotide triphosphates (dNTPs) are added as solutions of the salts of dATP, dCTP, dGTP, dUTP, and dTTP, such as disodium or lithium salts. In the present methods, a final concentration in the range of 1 μM to 2 mM each is suitable, and 100-600 μM is preferable, although the optimal concentration of the nucleotides may vary in the reverse transcription reaction depending on the total dNTP and divalent metal ion concentration, and on the buffer, salts, particular primers, and template. For longer products, i.e., greater than 1500 bp, 500 μM each dNTP may be preferred when using a Tris-HCl buffer.

20 dNTPs chelate divalent cations, therefore amount of divalent cations used may need to be changed according to the dNTP concentration in the reaction. Excessive amount of dNTPs (e.g., larger than 1.5 mM) can increase the error rate and possibly inhibits DNA polymerases. Lowering the dNTP (e.g., to 10-50 μM) may therefore reduce error rate. PCR reaction for amplifying larger size template may need more dNTPs.

25 One suitable buffering agent is Tris-HCl, preferably pH 8.3, although the pH may be in the range 8.0-8.8. The Tris-HCl concentration is from 5-250 mM, although 10-100 mM is most preferred. A preferred buffering agent is Bicine-KOH, preferably pH 8.3, although pH may be in the range 7.8-8.7. Bicine acts both as a pH buffer and as a metal buffer.

30 PCR is a very powerful tool for DNA amplification therefore very little template DNA is needed. However, in some embodiments, to reduce the likelihood of error, a higher DNA concentration may be used, though too many templates may increase the amount of contaminants and reduce efficiency.

5 Usually, up to 3 μ M of primers may be used, but high primer to template ratio can results in non-specific amplification and primer-dimer formation. Therefore it is usually necessary to check primer sequences to avoid primer-dimer formation. In a preferred embodiment, 0.1-0.5 μ M of primers are used.

3. Cycling Parameters

10 Denaturation time may be increased if template GC content is high. Higher annealing temperature may be needed for primers with high GC content or longer primers. Gradient PCR is a useful way of determining the annealing temperature. Extension time should be extended for larger PCR product amplifications. However, extension time may need to be reduced whenever possible to limit damage to enzyme.

15 The number of cycle can be increased if the number of template DNA is very low, and decreased if high amount of template DNA is used.

4. PCR Enhancing Factors And Additives

20 PCR enhancing factors may also be used to improve efficiency of the amplification. As used herein, a "PCR enhancing factor" or a "Polymerase Enhancing Factor" (PEF) refers to a complex or protein possessing polynucleotide polymerase enhancing activity (Hogrefe et al., 1997, Strategies 10::93-96; and U.S. Patent No. 6,183,997, both of which are hereby incorporated by references). For Pfu DNA polymerase, PEF comprises either P45 in native form (as a complex of P50 and P45) or as a recombinant protein. In the native complex of Pfu P50 and P45, only P45 exhibits PCR enhancing activity. The P50 protein is similar in structure to a bacterial flavoprotein. The P45 protein is similar in structure to dCTP deaminase and dUTPase, but it functions only as a dUTPase converting dUTP to dUMP and pyrophosphate. PEF, according to the present invention, can also be selected from the group consisting of: an isolated or purified naturally occurring polymerase enhancing protein obtained from an archeabacteria source (e.g., *Pyrococcus furiosus*); a wholly or partially synthetic protein having the same amino acid sequence as Pfu P45, or analogs thereof possessing polymerase enhancing activity; polymerase-enhancing mixtures of one or more of said naturally occurring or wholly or partially synthetic proteins; polymerase-enhancing protein complexes of one or more of said naturally

5 occurring or wholly or partially synthetic proteins; or polymerase-enhancing partially purified cell extracts containing one or more of said naturally occurring proteins (U.S. Patent No. 6,183,997, supra). The PCR enhancing activity of PEF is defined by means well known in the art. The unit definition for PEF is based on the dUTPase activity of PEF (P45), which is determined by monitoring the production of pyrophosphate (PPi) from dUTP. For example, PEF
10 is incubated with dUTP (10mM dUTP in 1x cloned Pfu PCR buffer) during which time PEF hydrolyzes dUTP to dUMP and PPi. The amount of PPi formed is quantitated using a coupled enzymatic assay system that is commercially available from Sigma (#P7275). One unit of activity is functionally defined as 4.0 nmole of PPi formed per hour (at 85°C).

Other PCR additives may also affect the accuracy and specificity of PCR reaction. EDTA less than 0.5 mM may be present in the amplification reaction mix. Detergents such as Tween-20TM and NonidetTM P-40 are present in the enzyme dilution buffers. A final concentration of non-ionic detergent approximately 0.1% or less is appropriate, however, 0.01-0.05% is preferred and will not interfere with polymerase activity. Similarly, glycerol is often present in enzyme preparations and is generally diluted to a concentration of 1-20% in the reaction mix. Glycerol (5-10%), formamide (1-5%) or DMSO (2-10%) can be added in PCR for template DNA with high GC content or long length (e.g., > 1kb). These additives change the T_m (melting temperature) of primer-template hybridization reaction and the thermostability of polymerase enzyme. BSA (up to 0.8 µg/µl) can improve efficiency of PCR reaction. Betaine (0.5-2M) is also useful for PCR over high GC content and long fragments of DNA.
25 Tetramethylammonium chloride (TMAC, >50mM), Tetraethylammonium chloride (TEAC), and Trimethylamine N-oxide (TMAO) may also be used. Test PCR reactions may be performed to determine optimum concentration of each additive mentioned above.

Various specific PCR amplification applications are available in the art (for reviews, see for example, Erlich, 1999, Rev Immunogenet., 1:127-34; Prediger 2001, Methods Mol. Biol.
30 160:49-63; Jurecic et al., 2000, Curr. Opin. Microbiol. 3:316-21; Triglia, 2000, Methods Mol. Biol. 130:79-83; MacClelland et al., 1994, PCR Methods Appl. 4:S66-81; Abramson and Myers, 1993, Current Opinion in Biotechnology 4:41-47; each of which is incorporated herein by references).

The subject invention can be used in PCR applications include, but are not limited to, i) hot-start PCR which reduces non-specific amplification; ii) touch-down PCR which starts at high annealing temperature, then decreases annealing temperature in steps to reduce non-specific PCR product; iii) nested PCR which synthesizes more reliable product using an outer set of primers and an inner set of primers; iv) inverse PCR for amplification of regions flanking a known sequence. In this method, DNA is digested, the desired fragment is circularized by ligation, then PCR using primer complementary to the known sequence extending outwards; v) AP-PCR (arbitrary primed)/RAPD (random amplified polymorphic DNA). These methods create genomic fingerprints from species with little-known target sequences by amplifying using arbitrary oligonucleotides; vi) RT-PCR which uses RNA-directed DNA polymerase (e.g., reverse transcriptase) to synthesize cDNAs which is then used for PCR. This method is extremely sensitive for detecting the expression of a specific sequence in a tissue or cells. It may also be use to quantify mRNA transcripts; vii) RACE (rapid amplification of cDNA ends). This is used where information about DNA/protein sequence is limited. The method amplifies 3' or 5' ends of cDNAs generating fragments of cDNA with only one specific primer each (plus one adaptor primer). Overlapping RACE products can then be combined to produce full length cDNA; viii) DD-PCR (differential display PCR) which is used to identify differentially expressed genes in different tissues. First step in DD-PCR involves RT-PCR, then amplification is performed using short, intentionally nonspecific primers; ix) Multiplex-PCR in which two or more unique targets of DNA sequences in the same specimen are amplified simultaneously. One DNA sequence can be use as control to verify the quality of PCR; x) Q/C-PCR (Quantitative comparative) which uses an internal control DNA sequence (but of different size) which compete with the target DNA (competitive PCR) for the same set of primers; xi) Recursive PCR which is used to synthesize genes. Oligonucleotides used in this method are complementary to stretches of a gene (>80 bases), alternately to the sense and to the antisense strands with ends overlapping (~20 bases); xii) Asymmetric PCR; xiii) In Situ PCR; xiv) Site-directed PCR Mutagenesis.

It should be understood that this invention is not limited to any particular amplification system. As other systems are developed, those systems may benefit by practice of this invention. A recent survey of amplification systems was published in.

5 B. Applications In Reverse Transcription

The term “reverse transcriptase” describes a class of polymerase characterized as RNA-dependent DNA polymerases. All known reverse transcriptases require a primer to synthesize a DNA transcript from an RNA template. Historically, reverse transcriptase has been used primarily to transcribe mRNA into cDNA which can then be cloned into a vector for further
10 manipulation (e.g., PCR amplification by a DNA-dependent DNA polymerase).

Avian myoblastosis virus (AMV) reverse transcriptase was the first widely used RNA-dependent DNA polymerase (Verma, 1977, *Biochem.Biophys.Acta* 473:1). The enzyme has 5'-3' RNA-directed DNA polymerase activity, 5'-3' DNA-directed DNA polymerase activity, and RNase H activity. RNase H is a processive 5' and 3' ribonuclease specific for the RNA strand of RNA-DNA hybrids (Perbal, 1984, A Practical Guide to Molecular Cloning, Wiley & Sons New
15 York). Errors in transcription cannot be corrected by reverse transcriptase because known viral reverse transcriptases lack the 3'-5' exonuclease activity necessary for proofreading (Saunders and Saunders, 1987, Microbial Genetics Applied to Biotechnology, Croom Helm, London). The use of the second enzyme in the subject composition provides proofreading for the reverse transcription reaction. A detailed study of the activity of AMV reverse transcriptase and its
20 associated RNase H activity has been presented by Berger et al., 1983, *Biochemistry* 22:2365-2372.

The reaction mixture for reverse transcription usually includes enzymes, aqueous buffers, salts, oligonucleotide primers, target polynucleotide, and nucleoside triphosphates. Depending
25 upon the context, the mixture can be either a complete or incomplete reverse transcription reaction mixture. The reaction mixture can be modified according to the conditions required by the second enzyme of the subject composition. It is known that cDNAs can be obtained from mRNAs in vitro using a reverse transcriptase (RNA-dependent DNA polymerase). The full length cDNA strands produced in turn may be used as a template for subsequent amplification
30 reaction (e.g., PCR) and the like.

Reverse transcription in combination with PCR (RT-PCT) is utilized to detect the presence of one or many specific RNA molecules which may be present in a sample. The method can be used to detect, for example, RNA from different organisms (such as viruses,

5 bacteria, fungi, plants, and animals), or RNA indicative of an infection, a disease state, or predisposition to a disease. For example, mRNA specific to tumor cells can be detected. The method is also useful for detecting a class of microorganisms or a group of related disease conditions.

Reverse transcription can generally be performed at any temperature within the
10 functional temperature range of the reverse transcriptase. Preferably, the temperature of incubation is any temperature at which the reverse transcriptase is functional and the primer remains hybridized to the RNA molecule. For non-thermostable reverse transcriptases, preferred temperatures are those temperatures that are at or around the optimum temperature for the reverse transcriptase. For most non-thermostable reverse transcriptases this temperature will be
15 between about 25°C and 45°C

U.S. Patent No. 5,994,079 discloses thermostable reverse transcriptases (herein incorporated by reference). Mn^{2+} is preferred as the divalent cation and is typically included as a salt, for example, manganese chloride ($MnCl_2$), manganese acetate ($Mn(OAc)_2$), or manganese sulfate ($MnSO_4$). If $MnCl_2$ is included in a reaction containing 10 mM Tris buffer, for example,
20 the $MnCl_2$ is generally present at a concentration of 0.5-7.0 mM; 0.8-1.4 mM is preferred when 200 μ M of each dGTP, dATP, dUTP, and, dCTP are utilized; an 1.2 mM $MnCl_2$ is most preferred.

A thermostable reverse transcriptase may retain at least 5% of its maximum activity at any temperature above 50°C or has an optimal temperature of at least 50°C. The highest
25 temperature at which a thermostable reverse transcriptase is functional can be quite high. For this reason, preferred temperature ranges for reverse transcription when a thermostable reverse transcriptase is used are most conveniently described in terms of the calculated melting temperature of a hybrid between the RNA molecule of interest and the primer. Such a melting temperature is referred to herein as the RNA/primer melting temperature (R/P T_m). Preferred
30 ranges include a temperature from 20°C below the melting temperature of a hybrid between the RNA molecule of interest and the primer and 5°C above the melting temperature of a hybrid between the RNA molecule of interest and the primer. In general, the closer the temperature is to the R/P T_m , the greater the degree of discrimination there will be between specific and non-

5 specific hybrids of the RNA and primer. If the temperature is close to the R/P T_m , however, decreased stability of specific hybrids may cause priming to be less efficient.

R/P T_m can be determined either by calculation or by empirical measurement. For calculating R/P T_m , any established formula for calculating stability of polynucleotide hybrids can be used. A preferred formula for calculating R/P T_m is $T_m = 81.5 + 16.6(\log M)^{+0.41}(\% G^+C) - 0.72(\% \text{ formamide})$, which was derived from studies on the stability of perfectly-matched
10 DNA:DNA hybrids. For RNA:DNA hybrids, incorporating formamide concentration in the formula does not hold because the relationship between formamide concentration and the depression of T_m is not linear. At 80% formamide, RNA:DNA hybrids are more stable than DNA:DNA hybrids, increasing the T_m by about 10 to 30°C depending on the sequence (Hames & Higgins, Polynucleotide Hybridisation: A Practical Approach (IRL Press Limited, Oxford, England. 1985)). Carrying out the reaction in 80% formamide can therefore also be used to suppress formation of DNA:DNA duplexes, to preferentially select RNA:DNA hybrids, and to estimate the T_m for R/P. Because the empirically derived formulas for the estimation of RNA:DNA hybrid T_m may not be as accurate for short DNA primers, the hybridization
15 temperature is preferably determined by assessing hybrid stability in 0.1-0.4 M monovalent cation at temperatures ranging from 40 to 60°C R/P T_m can also be determined empirically (Lesnick and Freier, 1995, Biochemistry 34:10807-10815, McGraw et al., 1990, Biotechniques 8:674-678; and Rychlik et al., 1990, Polynucleotides Res. 18:6409-6412).

The fidelity of viral reverse transcriptases, such as AMV-RT and MoMuLV-RT, may be
25 compared to thermoactive reverse transcriptases by a straightforward assay procedure described in U.S. Patent No. 5,994,079 (supra). Plasmid BS⁺ (Stratagene) can be used for such an assay. The plasmid encodes an α -complementing β -galactosidase activity and can be linearized with NdeI. T3 RNA polymerase is used to prepare a cRNA transcript of the α -donor region. After treatment of the cRNA with RNase-free DNase and isolation of the cRNA, the cRNA is used as a
30 template for a reverse transcription/amplification reaction. A reverse transcription primer complementary to the 3' end of the cDNA containing an NdeI sequence at its 5' terminus, and an upstream PCR primer comprising a PstI sequence at the 5' termini provide a 752 bp PCR product. The PCR product and the pBS⁺ vector are then digested with NdeI and PstI followed by ligation of the PCR product into the vector and transformation into a suitable host. The presence

5 of white colonies indicates that a mutation had occurred during the RT or PCR amplification. The assay provides means for assigning a relative value to the fidelity of the reverse transcriptase activity of various enzymes. Specific mutations can be determined by sequence analysis.

Following reverse transcription of RNA, the RNA can be removed from the RNA/cDNA hybrid by heat denaturation or by a number of other known means such as alkali, heat, or
10 enzyme treatment. Enzyme treatment may consist of, for example, treating the RNA/cDNA hybrid with RNase H. RNase H is specific for RNA strands within an RNA/DNA double-stranded molecule.

The subject composition is suitable for high fidelity transcribing and amplifying RNA from a number of sources. The RNA template may be contained within a polynucleotide preparation from an organism, for example, a viral or bacterial polynucleotide preparation. The
15 preparation may contain cell debris and other components, purified total RNA, or purified mRNA. The RNA template may be a population of heterogeneous RNA molecules in a sample or a specific target RNA molecule.

RNA suitable for use in the present methods may be contained in a biological sample suspected of containing a specific target RNA. The biological sample may be a heterogeneous
20 sample in which RNA is a small portion of the sample, as in for example, a blood sample or a biopsied tissue sample. Thus, the subject composition is useful for clinical detection and diagnosis. The RNA target may be indicative of a specific disease or infectious agent.

RNA may be prepared by any number of methods known in the art; the choice may
25 depend on the source of the sample and availability. Methods for preparing RNA are described in Davis et al., 1986, Basic Methods in Molecular Biology, Elsevier, NY, Chapter 11; Ausubel et al., 1987, Current Protocols in Molecular Biology, Chapter 4, John Wiley and Sons, NY; Kawasaki and Wang, 1989, PCR Technology, ed. Erlich, Stockton Press NY; Kawasaki, 1990, PCR Protocols: A Guide to Methods and Applications, Innis et al. eds. Academic Press, San
30 Diego; all of which are incorporated herein by references.

5 C. Detection Of Amplified Product

Detection of amplified polynucleotide product can be accomplished by any of a variety of well known techniques. In a preferred embodiment, the amplified product is separated on the basis of molecular weight by gel electrophoresis, and the separated products are then visualized by the use of polynucleotide specific stains which allow one to observe the discrete species of resolved amplified product present in the gel. Although numerous polynucleotide specific stains exist and would be suitable to visualize the electrophoretically separated polynucleotides, ethidium bromide is preferred.

Alternative methods suitable to detect the amplified polynucleotide product include hybridization-based detection means that use a labeled polynucleotide probe capable of hybridizing to the amplified product. Exemplary of such detection means include the Southern blot analysis, ribonuclease protection analysis using in vitro labeled polyribonucleotide probes, and similar methods for detecting polynucleotides having specific nucleotide sequences. See, for example, Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons, 1987.

Amplified products (e.g., by PCR or RT-PCR) using the subject composition of the invention can be used for subsequent analysis such as sequencing or cloning.

D. Application In Direct Cloning of PCR Amplified Product

While it is understood that the amplified product using subject composition can be cloned by any method known in the art. In one embodiment, the invention provides a composition which allows direct cloning of PCR amplified product.

The most common method for cloning PCR products involves incorporation of flanking restriction sites onto the ends of primer molecules. The PCR cycling is carried out and the amplified DNA is then purified, restricted with an appropriate endonuclease(s) and ligated to a compatible vector preparation.

A method for directly cloning PCR products eliminates the need for preparing primers having restriction recognition sequences and it would eliminate the need for a restriction step to

5 prepare the PCR product for cloning. Additionally, such method would preferably allow cloning PCR products directly without an intervening purification step.

U.S. Patent Nos. 5,827,657 and 5,487,993 (hereby incorporated by their entirety) discloses method for direct cloning of PCR products using a DNA polymerase which takes advantage of the single 3'-deoxy-adenosine monophosphate (dAMP) residues attached to the 3' termini of PCR generated nucleic acids. Vectors are prepared with recognition sequences that afford single 3'-terminal deoxy-thymidine monophosphate (dTMP) residues upon reaction with a suitable restriction enzyme. Thus, PCR generated copies of genes can be directly cloned into the vectors without need for preparing primers having suitable restriction sites therein.

Taq DNA polymerase exhibits terminal transferase activity that adds a single dATP to the 3' ends of PCR products in the absence of template. This activity is the basis for the TA cloning method in which PCR products amplified with Taq are directed ligated into vectors containing single 3'dT overhangs. Pfu DNA polymerase, on the other hand, lacks terminal transferase activity, and thus produces blunt-ended PCR products that are efficiently cloned into blunt-ended vectors.

In one embodiment, the subject invention comprises a Taq DNA polymerase as the first enzyme and a mutant Pfu DNA polymerase with reduced polymerization activity as the second enzyme. Taq DNA polymerase in the composition produces amplified DNA product with 3'-dAMP and allows direct cloning of the amplified product, while the mutant Pfu DNA polymerase provides fidelity for the amplification.

25 Kits

The invention herein also contemplates a kit format which comprises a package unit having one or more containers of the subject composition and in some embodiments including containers of various reagents used for polynucleotide synthesis, including synthesis in PCR. The kit may also contain one or more of the following items: polynucleotide precursors, primers, buffers, instructions, and controls. Kits may include containers of reagents mixed together in suitable proportions for performing the methods in accordance with the invention. Reagent

5 containers preferably contain reagents in unit quantities that obviate measuring steps when performing the subject methods.

EXAMPLES

The following examples are offered for the purpose of illustrating, not limiting, the subject invention.

10 **Example 1. Constructing Mutants Of Pfu DNA Polymerase With Reduced DNA Polymerase Activity**

15 We introduced mutations into Pfu DNA polymerase that were likely to reduce or eliminate DNA polymerase activity, while having minimal effects on proofreading activity. The mutations selected were identified from previous mutagenesis studies carried out using related Family B DNA polymerases. We made the same amino acid side chain substitutions in the polymerization domain at the following residues in Pfu (D405E, Y410F, T542P, D543G, K593T, Y595S) (Table 1).

Mutations were also introduced within the partitioning domain at amino acids 384-389 (SYTGGF) in Pfu DNA polymerase (Table 1).

20 The DNA template used for mutagenesis contained the Pfu *pol* gene, cloned into pBluescript (pF72 clone described in US 5,489,523) and expressed with an N-terminal His₆ tag for affinity purification. A modified QuikChange (Stratagene) protocol was used to insert the His₆ tag at the 5' end of the Pfu *pol* gene, just after the initiator ATG. The insertion reaction was carried out in two steps. In the first step, a standard QuikChange reaction was carried out in the presence of *Tth* ligase (10U/RXN) using only the His₆ forward primer. After 18 cycles, the reaction was *Dpn*I-digested for one hour at 37°C and then purified with the StrataPrep® Plasmid Miniprep Kit (Stratagene). The purified material served as the template in the second QuikChange reaction, which employed only the His₆ reverse primer. After 18 cycles, the second reaction was *Dpn*I-digested for one hour at 37°C, and then transformed. The His₆-Pfu *pol* construct was confirmed by both PCR amplification and sequencing using the Big Dye sequencing kit.

25
30

5 Point mutations were introduced into the *Pfu pol* gene using the QuikChange Site-Directed Mutagenesis Kit (Stratagene). Clones were sequenced to verify incorporation of the desired mutations.

Table 1: Activity of partially-purified His-tagged *Pfu* mutants (Nickel-resin eluates):

Mutation	Polymerase activity		Exonuclease activity		Rel. exo/pol vs. wt (1.0) ^s
	Cpms @ 50ng (500ng)	% wild type @ 50ng*	Cpms (50 ng)	% wild type [@]	

Partitioning					
S384G	46920	71	1425	≥100	2.3
S384K	66545	100	554	63	0.6
Y385N	1123	2	158	18	10.6
Y385W	10515 (24519)	16	36	4	0.3
Y385L	2383	4	180	21	5.7
Y385H	4276	6	91	10	1.6
Y385Q	386 (5431)	0.6	252	29	49.2
Y385S	1095 (4206)	2	578	66	39.8
Y385F	80685 (21580)	100	1008	≥100	0.9
T386E	48296	73	263	30	0.4
T386Y	47318	72	1112	≥100	1.8
T386G	46289	70	1011	≥100	1.6
G387S	648	1	169	19	19.7
G387P	258 (66)	0.4	500	57	146.2
G388A	2560	4	73	.008	2.2
G388S	74551	100	670	76	0.7
G388P	1222	2	202	23	12.5

F389Y	43455 (29809)	66	37	4	.06
F389L	72647	100	1054	≥100	1.1
F389V	30641	46	614	70	1.5
F389S	17998	27	1335	≥100	5.6
F389H	19623	30	543	62	2.1
	Polymerase activity		Exo/pol activity		
Polymerase	Cpms @ 5ng	% wild type @ 5ng [#]	Cpms exonuclease	Cpms Polymerase	Rel. exo/pol vs. wt (1.0) ^{&}
DXXSLYP					
D405E	69 (500ng)	<0.2	321	0	>396
Y410F	10181	27	698	16189	5.3
YXDTDS					
T542P	27	.07	1105	0	>1364
D543G	10	.03	704	687	127
T542P/D543G	23	.06	505	0	>623
KXY					
K593T	155	.4	668	0	>825
Y595S	6107	16	1072	2684	49

5 100% for wt *Pfu* equals: *66146 cpms; [#]38014 cpms; @877cpms
exo/pol for wt *Pfu* equals: ^{\$}0.01326; [&]0.0081

5 **Example 2. Affinity Purification Of His-Tagged Pfu DNA Polymerase Mutants**

Bacterial expression of Pfu mutants. Plasmid DNA was purified with the StrataPrep® Plasmid Miniprep Kit (Stratagene), and used to transform XL-10 Gold cells. Ampicillin resistant colonies were grown up in 1-5 liters of LB media containing Turbo Amp™ antibiotic (100µg/µl) at 37°C with moderate aeration. The cells were collected by centrifugation and stored at -20°C.

10 **Purification (His₆ tag protocol/batch binding method):** Cells pellets were resuspended in native binding buffer (20mM phosphate (pH 7.8), 500mM NaCl). Egg white lysozyme (100µg/ml) was added and the cells were incubated for 15 minutes on ice. Cell suspensions were subjected to sonication three times with a Bronson Sonifier 250 at a duty cycle of 80% and an output level of 5 for 45 seconds. The suspensions were left on ice to cool between sonication events. The lysate was cleared by centrifugation at 26,890g. The cleared lysates were added to 5mls of ProBond Ni resin (Invitrogen), equilibrated in native binding buffer, and the slurry was incubated for two hours with gentle agitation at 4°C. The resin was settled by low speed centrifugation (800Xg). The resin was washed three times with 4ml of native binding buffer (pH 7.8) by resuspending the resin, rocking the slurry for two minutes, and then separating the resin from the supernatant by gravity centrifugation. The resin was then washed in the same fashion with native wash buffer (20mM phosphate (pH 6.0), 500mM NaCl). Protein was eluted with two 5-ml additions of 350mM Imidazole elution buffer (20mM phosphate, 500mM NaCl, 350mM Imidazole (pH 6.0)) by resuspending the resin, rocking the slurry for five minutes, and then separating the resin from the supernatant by gravity centrifugation. Eluted proteins were spin concentrated using Centricon 30 centrifugal filter devices (Amicon). Protein samples were evaluated for size and purity by SDS-PAGE using Tris-Glycine 4-20% acrylamide gradient gels. Gels were stained with silver stain or Sypro Orange (Molecular Probes).

25 **Alternative expresssion/purification:** Alternatively, Pfu mutants were subcloned into the pCAL-n-EK vector (Affinity™ Protein Expression and Purification System) which contains an upstream, in-frame calmodulin binding peptide (CBP) tag for purifying fusion proteins with calmodulin agarose. Plasmid DNA was purified with the StrataPrep® Plasmid Miniprep Kit (Stratagene), and used to transform BL21(DE3) CodonPlus® cells. Ampicillin resistant colonies were grown up in 1-5 liters of LB media containing Turbo Amp™ antibiotic (100µg/µl) at 30°C

5 with moderate aeration. When cultures reached an absorbance at OD₆₀₀ of 0.6 to 1.0, the cells were induced with 1mM IPTG and incubated in the same manner for 2 hours to overnight (16 hours). The cells were collected by centrifugation and stored at -20°C.

Cells pellets were resuspended to an approximate concentration of 0.25g/ml in buffers identical or similar to calcium binding buffer (50mM Tris-HCL (pH 8.0), 150 mM NaCl, 1mM
10 magnesium acetate and 2mM CaCl). Egg white lysozyme (100µg/ml) was added and the cells were incubated for 15 minutes on ice. Cell suspensions were subjected to sonication three times with a Bronson Sonifier 250 at a duty cycle of 80% and an output level of 5 for 45 seconds. The suspensions were left on ice to cool between sonication events. The lysate was cleared by centrifugation at 26,890g.

15 The cleared lysates were added to 1ml of calmodulin agarose (CAM agarose), equilibrated in buffer, and the slurry was incubated with gentle agitation at 4°C. After two hours the reactions were centrifuged at 3000g for 5 minutes to collect the CAM agarose and recombinant protein. The lysate supernatant was removed and the CAM agarose was washed at least once by resuspending the resin in 50ml of calcium binding buffer followed by collection of
20 the CAM agarose by centrifugation as described above. The CAM agarose was transferred to a disposable 15ml column, packed, then washed with at least 200ml of calcium binding buffer. Recombinant proteins were eluted from the column by using a buffer similar or identical to 50mM Tris-HCl (pH 8.0), 1M NaCl, 2mM EGTA.

Protein samples were evaluated for size and purity by SDS-PAGE using Tris-Glycine 4-
25 20% acrylamide gradient gels. Gels were stained with silver stain or Sypro Orange (Molecular Probes).

Example 3. Assaying DNA Polymerase And 3'-5' Exonuclease Activities Of Pfu DNA Polymerase Mutants

Pfu mutant preparations were assayed for DNA polymerase and 3'-5' exonuclease
30 activities as follows.

DNA polymerase. DNA polymerase activity was measured by monitoring incorporation of radiolabelled TTP into activated calf thymus DNA. A suitable DNA polymerase reaction

5 cocktail contained: 1x PCR reaction buffer, 200 μ M each dATP, dCTP, and dGTP, 195 μ M TTP, 5 μ M [3 H]TTP (NEN #NET-221H, 20.5Ci/mmol; partially evaporated to remove EtOH), and 250 μ g/ml of activated calf thymus DNA (e.g., Pharmacia #27-4575-01). DNA polymerases (wt Pfu or Pfu mutants) were diluted in Pfu storage buffer and 1 μ l of each enzyme dilution was added to 10 μ l aliquots of polymerase cocktail. Polymerization reactions were conducted in
10 duplicate or triplicate for 30 minutes at 72°C. The extension reactions were quenched on ice, and then 5 μ l aliquots were spotted immediately onto DE81 ion-exchange filters (2.3cm; Whatman #3658323). Unincorporated [3 H]TTP was removed by 6 washes with 2xSCC (0.3M NaCl, 30mM sodium citrate, pH 7.0), followed by a brief wash with 100% ethanol. Incorporated radioactivity was measured by scintillation counting.

15 Reactions that lack enzyme were set up along with sample incubations to determine “total cpms” (omit filter wash steps) and “minimum cpms”(wash filters as above). Sample cpms were subtracted by minimum cpms to determine “corrected cpms” for each DNA polymerase.

To determine percent (%) activity relative to wild type Pfu, ~50-500ng of purified Pfu mutants were assayed in a nucleotide incorporation assay, alongside wild type Pfu diluted serially over the linear range of the assay (50-500pg; 0.003-0.03U).
20

Exonuclease assays. Exonuclease reactions were performed (in triplicate) by adding 4 μ l aliquots of diluted DNA polymerases (0.25-10U wt Pfu; 5-200ng) to 46 μ l of reaction cocktail. Reactions were incubated for 1 hour at 72°C. Reactions lacking DNA polymerase were also set up along with sample incubations to determine “total cpms” (no TCA precipitation) and
25 “minimum cpms” (TCA precipitation, see below).

Exonuclease reactions were stopped by transferring the tubes to ice. Sonicated salmon sperm DNA (150 μ l; 2.5 mg/ml stock) and TCA (200 μ l; 10% stock) were added to all but the “total cpms” tubes. The precipitation reactions were incubated for \geq 15 minutes on ice, and then spun in a microcentrifuge at 14,000rpm for 10 minutes. 200 μ l of the supernatant was removed,
30 being careful not to disturb the pellet, and transferred to scintillation fluid (Bio-Safe IITM, Research Products International Corp.). The samples were thoroughly mixed by inversion and then counted in a scintillation counter.

5 To determine percent (%) exonuclease activity relative to wild type Pfu, equivalent amounts of Pfu and purified Pfu mutants (which fall in the linear range of the assay; ~5-200ng Pfu) are assayed in an exonuclease assay.

10 **Results:** Several Pfu mutants exhibited reductions in DNA polymerase activity compared to wild type Pfu, when tested as partially purified (~50% purity) preparations eluted from nickel resins (Table 1). Pfu mutants showing <10% DNA polymerase activity and at least 10% exonuclease activity include the partitioning domain mutants: Y385QSNLH, G387SP, and G388P and the polymerase domain mutants: D405E, T542P, D543G, and K593T. The initial measurements of % DNA polymerase activity shown in Table 1 was considered as approximate estimates, due to the purity of the protein samples tested and uncertainties as to whether all protein amounts tested were in the linear range of the assay.

15 **Example 4. Purification Of Pfu DNA Polymerase Mutants By Conventional Column Chromatography**

20 The Pfu K593T and G387P mutants were purified as follows. Cells pellets (12-24 grams) were resuspended in 3 volumes of lysis buffer (buffer A: 50mM Tris HCl (pH 8.2), 1mM EDTA, and 10mM βME). Lysozyme (1 mg/g cells) and PMSF (1mM) were added and the cells were lysed for 1 hour at 4°C. The cell mixture was sonicated, and the debris removed by centrifugation at 15,000 rpm for 30 minutes (4°C). Tween 20 and Igelpal CA-630 were added to final concentrations of 0.1% and the supernatant was heated at 72°C for 10 minutes. Heat denatured *E. coli* proteins were then removed by centrifugation at 15,000 rpm for 30 minutes (4°C).

25 The supernatant was chromatographed on a Q-Sepharose™ Fast Flow column (~5ml column), equilibrated in buffer B (buffer A plus 0.1% (v/v) Igelpal CA-630, and 0.1% (v/v) Tween 20). Flow-through fractions were collected and then loaded directly onto a P11 Phosphocellulose column (1.6 x 10cm), equilibrated in buffer C (same as buffer B, except pH 7.5). The column was washed and then eluted with a 0-0.7M KCl gradient/Buffer C. Fractions containing Pfu DNA polymerase mutants (95kD by SDS-PAGE) were dialyzed overnight against buffer D (50mM Tris HCl (pH 7.5), 5mM βME, 5% (v/v) glycerol, 0.2% (v/v) Igelpal CA-630,

5 0.2% (v/v) Tween 20, and 0.5M NaCl) and then applied to a Hydroxyapatite column (1.0 x 1.3 cm; ~1ml), equilibrated in buffer D. The column was washed and Pfu DNA polymerase mutants were eluted with buffer D2 containing 400 mM KPO₄, (pH 7.5), 5mM βME, 5% (v/v) glycerol, 0.2% (v/v) Igepal CA-630, 0.2% (v/v) Tween 20, and 0.5 M NaCl. Purified proteins were spin concentrated using Centricon YM30 devices, and exchanged into Pfu final dialysis buffer
10 (50mM Tris-HCl (pH 8.2), 0.1mM EDTA, 1mM dithiothreitol (DTT), 50% (v/v) glycerol, 0.1% (v/v) Igepal CA-630, and 0.1% (v/v) Tween 20).

Results: His-tagged Pfu G387P and K593T mutants were purified by ion exchange/hydroxyappetite (IE/HA) chromatography. The purified protein preps were analyzed by SDS-PAGE and determined to be of ≥95% purity. The IE/HA purified mutants were tested in a nucleotide incorporation assay to more precisely quantify percent remaining DNA polymerase activity. As shown in Table 3, the Pfu G387P mutant exhibits no significant DNA polymerase activity (<100 cpms above background) when 0.2μg to 1.2μg of protein was assayed. These results indicate that the Pfu G387P mutant exhibits <0.01% ($<100/16661 \times 25/1200 = 0.00013$) of the DNA polymerase activity exhibited by wild type Pfu DNA polymerase. In comparison, the Pfu K593T mutant retains approximately 1-2% of the DNA polymerase activity of wild type Pfu.

Table 3. Residual Polymerase Activity in IE/HA Purified *Pfu* Mutant Preps:

<i>Pfu</i> DNA Polymerase	Amount. Assayed (ng)	Corrected cpms	Relative (%) Polymerase Activity	Mean Relative Polymerase Activity
Wild type	25	16,661	100	100
G387P	240	42	0.026	Cpms not significantly (<100cpms) above background; therefore, assume $<100/16661 \times 25/1200 = <0.01\%$
	600	0	-	
	1200	16	0.002	

K593T	80	1228	2.3	1.8
	200	1774	1.3	

5 **Example 5. Verifying The Presence Of Proofreading Activity In Pfu Mutants Under PCR Conditions**

A qualitative assay was used to verify that Pfu mutants retained 3'-5' exonuclease activity under PCR conditions. In this assay, the 900bp H α 1AT target is amplified with exo⁻ Pfu DNA polymerase (2.5U/50 μ l) using a forward primer containing a 3'dG, which produces a dG/dG mismatch upon annealing to the DNA template. The amplicon is amplified from human genomic DNA using the forward primer: 5'-GAG.GAG.AGC.AGG.AAA.GGT.GGA.AG-3' (100ng/50 μ l rxn) and the reverse primer: 5'-GAG.GTA.CAG.GGT.TGA.GGC.TACT.G-3' (100ng/50 μ l rxn). Amplification is carried out in the absence or presence of varying amounts of Pfu mutants (200ng to 3.6 μ g) on a Perkin/Elmer 9600 thermal cycler with the following program: (1 cycle) 95°C for 2.5 minutes; (30 cycles) 95°C for 40 seconds, 61°C for 10 seconds, 72°C for 2.5 minutes; (1 cycle) 72°C for 7 minutes. In the absence of proofreading activity, exo⁻ Pfu produces low yields of product, presumably because the enzyme can not efficiently extend a dG/dG mismatch. In the presence of Pfu mutants with proofreading activity, the 3'dG should be excised from the primer, thereby allowing exo⁻ Pfu to amplify the target in high yields. This PCR assay was used to verify that Pfu mutants tested in fidelity assays retained sufficient proofreading activity under PCR conditions to excise mismatched PCR primers. Moreover, the assay allowed us to determine the range of protein concentrations that could be added to PCR reactions without inhibition of amplification.

Results: As shown in Figure 1, amplifications conducted with exo⁻ Pfu alone produced low yields of product due to poor extension of the dG/dG mismatch. Product yields were significantly higher in the presence of the Pfu G387P and K593T mutants (125-500ng), presumably because these mutants excise the 3'dG from the primer, thereby allowing exo⁻ Pfu to efficiently amplify the target. Additional experiments showed that the polymerase deficient Pfu G387P and K593T mutants were unable to amplify the target in the absence of exo⁻ Pfu (or wild type Pfu).

5 **Example 6. Range Of Ratios Of Exonuclease And Polymerase Activities To Use In Blends (Table 4)**

Enzyme blend					
Polymerase proficient enzyme		Polymerase deficient enzyme			
Polymerase	Amount Polymerase (3'-5' Exo)	Pfu Mutant	Range of Amounts Tested that Produce Highest Fidelity and Yield		
			Ng	Polymerase (U) [@]	3'-5' Exo (U) [@]
Pfu/ Pfu <i>Turbo</i> *	2.5-5U (0.05-.09U exo)	G387P	600 ^{&}	<0.003	0.56 ^{&}
			1800	<0.009	1.68
			3600	<0.018	3.36
Taq [#]	2.5U (0)	G387P	600 ^{&}	<0.003	0.56 ^{&}
			3600 ^s	<0.018	3.36 ^s

*600ng-3600ng of Pfu G387P provided similar improvements in the accuracy of Pfu; 1800ng was the highest amount that could be added with no inhibition of any PCR reaction tested (up to 17kb); at least one PCR system amplified successfully in the presence of 3600ng Pfu G387P

- 10 [#]600ng of Pfu G387P increased accuracy of Taq by 5-fold, while 3600ng increased accuracy by 8-fold; several targets <5kb successfully amplified in the presence of 3600ng of Pfu G387P; maximum amount of Pfu G387P that can be used in PCR may be >3600ng.

[&]minimum amount could be lower; haven't tested amounts <600ng in a fidelity assay

^smaximum amount could be higher; haven't tested amounts >3600ng in PCR

- 15 [@]calculations assume 50-100ng of Pfu equals 2.5U DNA polymerase activity (specific activity is 2.6×10^3 U/mg; measured at 72°C using activated calf thymus DNA in a universal assay buffer (50mM Tris-HCl (pH 8.0), 5mM MgCl₂, 1mM DTT, and 50µg/ml BSA)) and that the specific

5 exonuclease activity of Pfu is 935U/mg (measured at 72° in the absence of dNTPs using double-stranded *Pal* I-digested λ DNA in a universal assay buffer (70mM Tris HCl (pH 8.8), 2mM MgCl₂, 0.1% Triton-X, and 100μg/ml BSA)).

Example 7. PCR Amplification With Pfu Or Taq DNA Polymerase Blends Containing Pfu Mutants

10 **Pfu blends.** PCR reactions were conducted under standard conditions in cloned Pfu PCR buffer (10mM KCl, 10mM (NH₄)₂SO₄, 20mM Tris HCl (pH 8.8), 2mM Mg SO₄, 0.1% Triton X-100, and 100μg/ml BSA) with 2.5-5U *PfuTurbo* DNA polymerase (2.5U/μl cloned Pfu DNA polymerase plus 1U/μl native or 2U/μl cloned *Pyrococcus furiosus* dUTPase (PEF)) and varying concentrations of polymerase deficient Pfu mutants (e.g., 0.2 to 3.6μg). For genomic targets 0.3-15 9kb in length, PCR reactions contained 2.5U *PfuTurbo* DNA polymerase, 100ng of human genomic DNA, 200μM each dNTP, and 100ng of each primer. For genomic targets 11.9kb and 17kb in length, PCR reactions contained 5U *PfuTurbo* DNA polymerase, 250ng of human genomic DNA, 500μM each dNTP, and 200ng of each primer.

20 **Taq blends.** PCR reactions were conducted under standard conditions in Herculanase PCR buffer (50mM Tricine (pH 9.1), 8mM (NH₄)₂SO₄, 2.3mM MgCl₂, 0.1% Tween-20, and 75μg/ml BSA) with 2.5U cloned Taq DNA polymerase, 1U of native or 2U cloned *Pyrococcus furiosus* dUTPase (PEF)), and varying concentrations of polymerase deficient Pfu mutants.

Cycling Conditions (Table 5):

Target size (kb)	Target gene	Cycling Parameters
0.3	Aldolase B	(1 cycle) 95°C 2 min (30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 1 min (1 cycle) 72°C 7 min

0.9	H α 1AT	<p>(1 cycle) 95°C 2 min</p> <p>(30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 1 min</p> <p>(1 cycle) 72°C 7 min</p>
2.3	Pfu pol (5ng plasmid DNA)	<p>(1 cycle) 95°C 2 min</p> <p>(30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 3 min</p> <p>(1 cycle) 72°C 7 min</p>
2.6	H α 1AT	<p>(1 cycle) 95°C 2 min</p> <p>(30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 3 min</p> <p>(1 cycle) 72°C 7 min</p>
4	H α 1AT	<p>(1 cycle) 95°C 2 min</p> <p>(30 cycles) 95°C 40 sec, 54°C 30 sec, 72°C 5 min</p> <p>(1 cycle) 72°C 7 min</p>
9.3	H α 1AT	<p>(1 cycle) 95°C 2 min</p> <p>(30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 18 min</p> <p>(1 cycle) 72°C 10 min</p>

11.9	H α 1AT	(1 cycle) 95°C 2 min (30 cycles) 95°C 40 sec, 58°C 30 sec, 72°C 24 min (1 cycle) 72°C 10 min
17	β globin	(one cycle) 92°C 2 min (10 cycles) 92°C 10 sec, 63°C 30 sec, 68°C 30 min (20 cycles) 92°C 10 sec, 63°C 30 sec, 68°C 30 min (plus 10 sec/cycle) (one cycle) 68°C 10 min

Results (Pfu blend PCR performance): As shown in Figure 2, adding 600ng of the Pfu G387P mutant to Pfu (in the presence of PEF/dUTPase), has minimal effects on PCR product yield. Additional experiments have shown that up to 1800ng of the Pfu G387P mutant can be added without significantly reducing PCR product yield.

Results (Taq blend PCR performance): As shown in Figure 3, adding the Pfu G387P mutant to Taq, in the presence of PEF/dUTPase, significantly increases PCR product yields when amplifications are performed in a reaction buffer that supports the activity of both Taq and Pfu DNA polymerases. One such buffer is the Herculase PCR buffer, which was developed specifically for Herculase Enhanced DNA polymerase (3.33U/ μ l cloned Pfu, 1.67U/ μ l cloned Taq, 2U/ μ l cloned *Pyrococcus furiosus* dUTPase). In the example shown in Figure 3, a 4kb target could not be amplified in high yield using Taq alone in Taq, Pfu, or Herculase PCR buffer. In the presence of the Pfu G387P mutant (and dUTPase), the 4 kb target could be amplified in cloned Pfu buffer (moderate yield) but not Taq buffer, consistent with the buffer preferences of the Pfu G387P mutant. Other experiments have shown that the Pfu G387P mutant inhibits PCR reactions carried out with Taq in Taq PCR buffer, suggesting that the Pfu G387P mutant binds

5 the 3' ends of PCR products without excising mismatches and dissociating (due to inactivity in Taq buffer), and blocks further product extension. As expected, highest product yields are obtained with Taq plus Pfu G387P blends in the presence of Herculase buffer, since both enzymes are highly active in this particular buffer. The Pfu G387P mutant is thought to enhance the yields of Taq PCR reactions (in buffers where Pfu is active) by excising mispairs that would otherwise stall Taq.

Example 8. Measuring The Fidelity Of DNA Polymerase Blends Containing Pfu DNA Polymerase Mutants

15 The error rates of Pfu and Taq blends containing the Pfu G387P and K593T mutants were tested in the *lacI* PCR fidelity assay described in Cline, J., Braman, J.C., and Hogrefe, H.H. (96) NAR 24:3546-3551. Briefly, a 1.9kb fragment encoding the *lacIOlacZα* target gene was amplified from pPRIAZ plasmid DNA using 2.5U PfuTurbo in cloned Pfu PCR buffer or 2.5U Taq in Taq or Herculase PCR buffer. Varying amounts of the Pfu G387P (600-3600ng) and K593T (200-1200ng) mutants were added to certain reactions. For comparative purposes, the *lacI* target was also amplified with Pfx (*Thermococcus* sp. KOD DNA polymerase; Invitrogen) and Tgo (*Thermococcus gorgonarius* DNA polymerase; Roche) using the manufacturers' recommended PCR buffer. The *lacI*-containing PCR products were then cloned into lambda GT10 arms, and the percentage of *lacI* mutants (MF, mutation frequency) was determined in a color screening assay, as described (Lundberg, K.S., Shoemaker, D.D., Adams, M.W.W., Short, J.M., Sorge, J.A., and Mathur, E.J. (1991) Gene 180:1-8). Error rates are expressed as mutation frequency per bp per duplication (MF/bp/d), where bp is the number of detectable sites in the *lacI* gene sequence (349) and d is the number of effective target doublings. For each enzyme, at least two independent PCR amplifications were performed.

30 Error rate measurements have shown that Pfu and PfuTurbo DNA polymerases exhibit an average error rate which is ~2-fold lower than that of Vent, Deep Vent, and Pfx (KOD) DNA polymerases, 3 to 6-fold lower than those of DNA polymerase mixtures, and 6- to 12-fold lower than that of Taq DNA polymerase.

Results (Pfu blend): As shown Table 6, adding 600ng-3600ng of the IE/HA-purified Pfu G387G mutant reduced the error rate of PfuTurbo DNA polymerase by 3.2 to 3.5-fold (assay 1)

and by 1.8 to 2.8-fold (assay 2) in two independent fidelity assays. Under these conditions, the Pfu G387G mutant was present at approximately a 12-fold (600ng; <0.003U polymerase activity; 0.56U exonuclease activity) to 72-fold (3600ng; <0.018U polymerase activity; 3.4U exonuclease activity) molar excess over wild type Pfu. These assumptions were based upon 2.5U Pfu equaling ~50ng protein, the Pfu G387G mutant exhibiting <0.01% DNA polymerase activity, and wt Pfu exhibiting a specific activity of 935U/mg). As discussed in Example 5, up to 1800ng of the IE/HA-purified Pfu G387G mutant can be added to PCR reactions without significantly reducing PCR product yield.

In comparison, adding 200ng of the Pfu K593T mutant reduced the error rate of PfuTurbo DNA polymerase slightly (40%), while the addition of 600ng and 1200ng (~12 to 24-fold excess over Pfu) increased error rate by 2.8- and 7.3-fold, respectively. At 600-1200ng, approximately 0.5-1U of additional DNA polymerase activity is added to the PCR reaction (Pfu K593T mutant exhibits 1-2% polymerase activity). The K593T mutation significantly increases the misincorporation or mispair extension rate of Pfu, and when added at high amounts (corresponding to ≥ 0.5 U), the Pfu K593T mutant dramatically increases the error rate of wild type Pfu.

Results (Taq blend): As shown Table 7, adding 600ng and 3600ng of the Pfu G387G mutant reduced the error rate of Taq DNA polymerase by 5.1- and 8.3-fold, respectively. Therefore, the error rate of Taq in the presence of the Pfu G387G mutant, can equal the error rate of Pfu alone.

Table 6. Fidelity of *Pfu* Blends Containing IE/HA Purified *Pfu* Mutants:

PCR Enzyme	His- <i>Pfu</i> Mutant	Mutant Amount		Error rate* ($\times 10^{-6}$)		Mean Relative Accuracy (<i>Pfu</i>)
		ng (\times <i>Pfu</i>)	pol units	Assay 1	Assay 2	
<i>Pfu</i>	none	-	-	5.55	3.60	1.0

	G387P	600 (12x)	<0.003	1.60	2.06	2.6
		1800 (36x)	<0.009	1.65	1.18	3.2
		2400 (48x)	<0.012	Nd	1.30	2.8
		3600 (72x)	<0.018	1.75	nd	3.2
	K593T	200 (4x)	0.18	3.9	nd	1.4
		600 (12x)	0.54	15.7	nd	0.4
		1200 (24x)	1.1	40.3	nd	0.1
<i>Tgo</i>	none	-	-	nd	6.10	0.6
<i>Taq</i>	none	-	-	34.7	19.0	0.2

*mean of duplicate measurements

Table 7. Fidelity of *Taq* Blends Containing IE/HA Purified *Pfu* Mutants:

PCR Enzyme	His- <i>Pfu</i> Mutant	Mutant Amount (ng)	Error rate* (x 10 ⁻⁶)	Relative Accuracy (<i>Pfu</i>)
<i>Taq</i>	none	-	34.7	0.16
	G387P	600	6.8	0.82
		3600	4.2	1.32
	K593T	200	37.0	0.15

<i>Pfu</i>	none	-	5.6	1.0
	G387P	600	1.60	3.47
		3600	1.75	3.17
	K593T	200	3.90	1.42

*mean of duplicate measurements

Example 9. Determining The TA Cloning Efficiencies Of PCR Products Amplified With Taq In The Presence Of Pfu Mutants

To determine the effects of polymerase deficient Pfu mutants on the terminal transferase activity of Taq, we amplified a series of amplicons with Taq in the absence of the Pfu G387P mutant (in Taq PCR buffer) or in the presence of the Pfu G387P mutant (in Herculanase PCR buffer). Similar amplifications were performed using Pfu*Turbo* and Herculanase in their recommended PCR buffers. PCR product yields were quantified by analyzing the products on 1% agarose gels, stained with SYBR gold. The same amount of each PCR product was added to 1µl of the pCR 2.1-TOPO vector (Invitrogen) in a final reaction volume of 6µl, according the manual for the TOPO TA Cloning Kit (#K4500-01). The reactions were incubated for 5 minutes at room temperature, and then transferred to ice. The reactions were transformed into One-Shot cells (Invitrogen), according to the manufacture's recommendations. Aliquots of each transformation were plated on ampicillin/IPTG/X-gal plates, prepared as described in the

Invitrogen TOPO TA Cloning manual. The frequency of clones containing the desired insert (% cloning efficiency) was quantified as the number of (white colonies)/(total number of colonies plated).

Results: As shown in Table 8, PCR products amplified with Taq in the presence of the Pfu G387P mutant are cloned into the TOPO TA cloning vector as efficiently as PCR products amplified with Taq alone. In contrast, PCR products amplified with Pfu*Turbo* DNA polymerase

- 5 are cloned into the TOPO TA cloning vector much less efficiently, presumably due to the lack of
3' dAs. As discussed in Example 7, PCR products amplified with Taq blends containing the Pfu
G387P mutant, should also exhibit fewer errors (5- to 8-fold less) compared to PCR products
amplified with Taq alone. Therefore, Taq blends containing the Pfu G387P mutant should be
useful to researchers using TA cloning methods, but desiring high-fidelity amplification of inserts.
- 10 The high TA cloning efficiencies obtained in the presence of the Pfu G387P mutant indicates
that 3' dAs added by Taq during PCR are unexpectedly resistant to exonucleolytic degradation.
Presumably, Pfu DNA polymerase is not very efficient at excising 3'dA residues from double-
stranded PCR products in the presence of nucleotides.

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

5 Table 8. TopoTA Cloning Efficiencies:

PCR Product (bp)	PCR enzyme/blend			Cloning efficiency (%)
	DNA polymerase	Pol' <i>Pfu</i> mutant		
		mutant	amount (μg)	
900	<i>Taq</i>	none	-	89
		G387P	0.6	80
		G387P	3.6	89
	<i>Pfu</i>	none	-	8
300	<i>Taq</i>	none	-	69
		G387P	0.6	73
		G387P	3.6	78
	<i>Pfu</i>	none	-	33
	Herculase	None	-	46
2300	<i>Taq</i>	None	-	83
		G387P	0.6	88
		G387P	3.6	92
	<i>Pfu</i>	None	-	22
	Herculase	None	-	85

OTHER EMBODIMENTS

The foregoing examples demonstrate experiments performed and contemplated by the present inventors in making and carrying out the invention. It is believed that these examples include a disclosure of techniques which serve to both apprise the art of the practice of the



5 invention and to demonstrate its usefulness. It will be appreciated by those of skill in the art that the techniques and embodiments disclosed herein are preferred embodiments only that in general numerous equivalent methods and techniques may be employed to achieve the same result.

All of the references identified hereinabove, are hereby expressly incorporated herein by reference to the extent that they describe, set forth, provide a basis for or enable compositions
10 and/or methods which may be important to the practice of one or more embodiments of the present inventions.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215